

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 002 801 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
18.06.2003 Bulletin 2003/25

(21) Application number: 99122865.1

(22) Date of filing: 17.11.1999

(51) Int Cl.7: **C07F 9/50**, C07F 9/655,
C07B 31/00, C07B 53/00,
C07C 25/18, C07C 33/46,
C07F 15/00, B01J 31/28,
C08F 112/08

(54) Chiral diphenyldiphosphines and d-8 metal complexes thereof

Chirale Diphenyldiphosphine und d-8 Metall-Komplexe davon

Diphenyldiphosphines chirales et leurs complexes métalliques d-8

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: 19.11.1998 CH 231998

(43) Date of publication of application:
24.05.2000 Bulletin 2000/21

(73) Proprietor: Solvias AG
4057 Basel (CH)

(72) Inventors:

- Pugin, Benoît
4142 Muenchenstein (CH)
- Steiner, Ivo
6005 Luzern (CH)
- Aufdenblatten, Rhony Niklaus
3920 Zermatt (CH)
- Togni, Antonio
8332 Russikon (CH)

(74) Representative:
Dannappel, Hans-Jochen, Dr. et al
Solvias AG,
Patente,
WKL-402.4.26,
Klybeckstrasse 191
4002 Basel (CH)

(56) References cited:

EP-A- 0 104 375 WO-A-93/15090
WO-A-98/01457 WO-A-98/12202

- DATABASE Xfire [Online] Beilstein Institut für Literatur der organischen Chemie XP002132240 & ORG. MAGN. RESON., vol. 9, 1977, page 44
- SCHMID R ET AL: "Axially dissymmetric bis(triaryl)phosphines in the biphenyl series: synthesis of (6,6'-dimethylbiphenyl-2,2'-diyl)bis(diphenylphosphine) (BIPHEMP) and analogs, and their use in rhodium(I) catalyzed asymmetric isomerizations of N,N-diethylnerylamine" HELV. CHIM. ACTA (HCACAV,0018019X);1988; VOL.71 (4); PP.897-929, XP002132237 F. Hoffmann-La Roche und Co. A.-G.;Basel; CH-4002; Switz. (CH)
- MCGINN F.A.: "Direct configurational intercorrelation of 6,6'-dinitro, 6,6'-dichloro-, and 6,6'-dimethyl-2,2'-diphenic acid." JOURNAL OF THE AMERICAN CHEMICAL SOCIETY., vol. 80, no. 2, 1958, pages 476-480, XP002132238 AMERICAN CHEMICAL SOCIETY, WASHINGTON, DC., US ISSN: 0002-7863
- SCHMID R ET AL: "New developments in enantioselective hydrogenation" PURE APPL. CHEM. (PACHAS,00334545);1996; VOL.68 (1); PP.131-8, XP000884387 F. Hoffmann-La Roche Ltd.;Process Res. Units; Basel; CH-4002; Switz. (CH)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 002 801 B1

Description

[0001] The present invention relates to mono- and di-(halogenmethyl)-diphenyldiiodides, -dibromides and -diamines; mono- and di-(hydroxymethyl)-diphenyldiiodides, -dibromides and -diamines and mono- and di-(aminomethyl)-diphenyldiiodides, -dibromides and -diamines; mono- and di-(hydroxymethyl)-diphenyldiphosphines and mono- and di-(aminomethyl)-diphenyldiphosphines, as well as 2-hydroxypropane-1,3-dioxyl-diphenyldiphosphines; mono- and di-(hydroxymethyl)-diphenyldiphosphines and mono- and di-(aminomethyl)-diphenyldiphosphines, as well as 2-hydroxypropane-1,3-dioxyl-diphenyldiphosphines, the hydroxyl groups or amino groups of which are provided with functional groups via a bridging group; inorganic and organic polymeric carriers which are immobilised with said diphosphines; metal complexes of the monomeric and immobilised diphosphines; and the use of the metal complexes as homogeneous and enantioselective catalysts in the synthesis of organic compounds, for example hydrogenation.

[0002] In Pure and Appl. Chem., Vol. 68, No. 1, pp. 131-138 (1996), R. Schmid et al. describe atropisomeric 6,6'-dimethyl- and 6,6'-dimethoxy-2,2'-diphenyldiphosphines as chiral ligands in metal complexes, which are used for the hydrogenation of prochiral ketones and olefins, whereby high optical yields may be attained. The catalysts can only be extracted from the reaction mixtures with difficulty and incompletely, so that it is impossible to reuse them for further reactions.

[0003] In WO 98/01457, B. Pugin et al. describe the functionalisation of chiral ferrocenyldiphosphines as ligands for metal complexes and the immobilisation on inorganic and organic carriers, which may be used as enantioselective hydrogenation catalysts. These catalysts may be easily separated from the reaction mixture and reused. In Organic Magnetic Resonance 9(1), 44-48 (1997) is disclosed 2,2'-bis-(bromomethyl)-6,6'-dinitrobiphenyl. In Helvetica Chemical Acta 71(4), 897-929 (1988) is described 2,2'-bis-(hydroxymethyl)-6,6'-dibromobiphenyl and in J.A.C.S. 80(2), 476-480 (1958) is disclosed 2,2'-bis-(hydroxymethyl)-6,6'-diaminobiphenyl. In EP-A-0 104 375 and WO 93/15090 are disclosed 2,2'-substituted 6,6'-diphosphindiphenyl ligands for the asymmetric catalysed hydrogenation of unsaturated compounds, whereby the 2,2'-Substituents are not functionalised.

[0004] It has now been found that, in a simple manner, functionalised 2,2'-diphenyldiphosphines can be prepared and may be immobilised both on inorganic and on organic polymeric carriers, and can also be used as water-soluble and/or extractable and/or adsorbable ligands/catalysts. The immobilised diphosphine ligands bond with d-8 metals such as rhodium, ruthenium and iridium complexes which may be used as highly effective catalysts in enantioselective hydrogenation of carbon-carbon, carbon-nitrogen or carbon-oxygen double bonds. The selectivity, activity and total yield for immobilised systems are surprisingly high.


[0005] The catalysts may be easily separated from the reaction solution and reused. Almost no metal or ligand losses occur. In addition, immobilised diphenyldiphosphine ligands especially on inorganic carriers have surprisingly high stability, which is especially important for reuse. Therefore, using these immobilised catalysts, large-scale hydrogenation may be carried out especially economically.

[0006] The reaction to be catalysed may be carried out even heterogeneously or homogeneously through the choice of polymer, for example in the case of polymer-bound diphosphine ligands. The polymer may be prepared in such a way, or also subsequently specifically modified in such a way that the polymer-bound catalyst dissolves in the reaction medium, and can be easily separated after the reaction by filtration, ultrafiltration, extraction or adsorption on carriers, and then reused. The catalysts can be reused several times. Through the choice of polymer, the catalyst may be optimally adapted to the reaction medium during the hydrogenation step, and then completely separated, which is important in particular for hydrogenation carried out on a large scale.

[0007] The production of these immobilised or extractable and/or adsorbable diphenyldiphosphines is made possible only by providing correspondingly functionalised diphenyldiphosphines. Therefore, particular importance is placed on these intermediates and their preparation.

[0008] In all cases, recovery of the noble metals contained therein is simplified if the catalyst has to be exchanged after frequent recycling. Frequently, further purification of the hydrogenated product can be dispensed with, since the catalyst can be removed practically quantitatively.

[0009] A first object of the invention is compounds of formula I,



(I),


wherein

R_1 is methyl chloride, methyl bromide or methyl iodide, R_2 is C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy or has the same significance as R_1 , and R_3 is Br, I or $-NH_2$.

R_1 is preferably methyl bromide. R_2 is preferably alkyl, and as alkyl preferably signifies ethyl and most preferably methyl. R_2 as alkoxy preferably signifies methoxy or ethoxy. R_3 is preferably Br or I.

[0010] The preparation thereof may be effected in known manner by the radical halogenation of the methyl groups of 2,2'-di- R_3 -6,6'-dimethyl-diphenyl with appropriate halogenation agents, for example Cl_2 , Br_2 , I_2 , interhalogen compounds such as $ClBr$, ClI , or $SOCl_2$, $SOBr_2$, SOI_2 , and organic halogen compounds such as CF_3Cl , CF_3Br , CF_3I , CCl_3I , as well as N-halogenated acid amides, for example N-chloro-, -bromo- and -iodosuccinimide. Depending on the amount of halogenation agent, mono- or dihalogen-methyl-diphenyls are primarily obtained, whereby mixtures of the compounds may be separated by distillation, by chromatographic methods or by crystallisation. The oily or crystalline compounds of formula I are valuable initial products for the production of atropisomeric diphosphines.

[0011] A further object of the invention is compounds of formula II,



(II),

wherein

R_3 is Br, I or $-NH_2$,


R_4 is hydroxymethyl, aminomethyl, hydroxy-, amino- or cyano- C_2 - C_8 -alkoxy, hydroxy-, amino- or cyano- C_2 - C_8 -alkoxymethyl, or hydroxy-, amino- or cyano- C_2 - C_8 -alkylaminomethyl, and R_5 has the same significance as R_4 , or R_5 is C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy, or

R_4 and R_5 together are $HOCH(CH_2-O-)_2$, $H_2NCH(CH_2-O-)_2$, or hydroxy-, amino- or cyano- C_2 - C_8 -alkyl $OCH(CH_2-O-)_2$, except 2,2'-Dihydroxymethyl-6,6'-dibromodiphenyl and 2,2'-Dihydroxymethyl-6,6'-diaminodiphenyl.

R_3 is preferably Br or I and especially I. C_2 - C_8 -alkyl in the hydroxy-, amino- or cyanoalkyl groups is preferably C_2 - C_6 -alkyl, more preferably C_2 - C_4 -alkyl, for example C_2 -, C_3 -alkyl or C_4 -alkyl. R_5 as C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy may be for example methyl, ethyl, propyl, butyl, methoxy, ethoxy, propyloxy and butyloxy; methyl and methoxy are preferred. If R_5 signifies alkyl, R_4 is preferably hydroxymethyl, aminomethyl, hydroxy-, amino- or cyano- C_2 - C_8 -alkoxymethyl, or hydroxy-, amino- or cyano- C_2 - C_8 -alkylaminomethyl. If R_5 signifies alkoxy, R_4 is preferably amino- or cyano- C_2 - C_8 -alkoxy. The hydroxy, amino or cyano groups are preferably primary groups.

[0012] The compounds of formula II may be produced by known synthesis methods. Thus, the methyl halide group in the compounds of formula I can be hydroxylated or aminated in known manner. These hydroxy or amino compounds, or the known 2-alkoxy-2'-hydroxy-6,6'-substituted diphenyl, or the known 2,2'-dihydroxy-6,6'-substituted diphenyl, or


a compound of formula



which is obtainable by reacting 2,2'-dihydroxy-6,6'-substituted diphenyl with epichlorohydrin or 1,3-dichloro-2-hydroxypropane, is reacted with cyanoalkenyl and then the cyano group is hydrogenated, or reacted with hydroxyhalogen- or aminohalogen-alkanes, or with ethylene oxide or aziridine. The HO group may be substituted by a NH₂ group in known manner, for example first of all halogenated and then reacted with 1,1'-carbonyl-diimidazole; hydrolysis then yields the free amine.

[0013] The compounds of formula II are eminently suitable for producing corresponding diphenyldiphosphines, in which the functional groups may undergo prior or subsequent modification in conventional manner by conversion into other functional groups or by a reaction with difunctional chain extenders, for example epoxides, hydroxyalkyl cyanates, halo-alkanols, halo-alkane nitriles, halo-alkane phthalimides, dicarboxylic acids or diisocyanates. The phosphine groups are introduced in known manner [see Pure and Appl. Chem., Vol. 68, No. 1, pp. 131-138 (1996)] by reacting compounds of formula II with lithium alkyls, for example lithium butyl, and reacting these with secondary phosphine halides, for example chlorides. The functional groups may be provided with appropriate protecting groups, a large number of which are known. The introduction of the phosphine groups may also take place in stages, whereupon unsymmetrically substituted diphenyldiphosphines are obtainable. This functionalisation is described in detail in WO 98/01457.

[0014] Further objects of the invention are also the compounds of formula III,




(III),

wherein

R₆ and R₇ signify identical or different secondary phosphino,
 R₈ is -CH₂-OH, -CH₂-NH₂, -CH₂-O-B-(FU)_p, -CH₂-NR'-B-(FU)_p, or -O-B-(FU)_p,
 R₉ has the same significance as R₈ or is C₁-C₄-alkyl or C₁-C₄-alkoxy, or
 R₈ and R₉ together signify HOCH(CH₂-O-)₂, H₂NCH(CH₂-O-)₂, (FU)_p-B-OCH(CH₂-O-)₂ or (FU)_p-B-R'NCH(CH₂-O-)₂,
 R' is H or C₁-C₄ alkyl;
 B is a bridging group,
 FU is a functional group,
 p is a number from 1 to 6, and
 NH₂ groups are present as such or as masked isocyanate groups.


[0015] In formula III, p is preferably a number from 1 to 4, most preferably 1 to 3. Preferred compounds having more than one functional group are those of formula IIIb



wherein R_6 and R_7 are defined as above, R_8 is C_1 - C_4 -alkyl and preferably methyl, R_{10} is C_2 - C_{12} , preferably C_2 - C_6 , most preferably C_2 to C_4 -alkylene, X'' is $-COOH$ and X' is $-O-$, $-NH-$ or $-N(C_1-C_4\text{-alkyl})$, as well as the amides, esters and salts thereof, especially alkali or alkaline earth metal salts.

[0016] If R_8 and R_9 contain a primary amino group, this may be converted by known methods into masked isocyanate groups, which similarly represent valuable functional groups.

[0017] The secondary phosphino may correspond to formula $-PR_{10}R_{11}$, wherein R_{10} and R_{11} , independently of one another, are C_1 - C_{12} -alkyl, C_5 - C_{12} -cycloalkyl; phenyl, C_5 - C_{12} -cycloalkyl substituted by C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy; or phenyl mono- to trisubstituted by C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, $-SiR_{12}R_{13}R_{14}$, halogen, $-SO_3M$, $-CO_2M$, $-PO_3M$, $-NR_{15}R_{16}$, $[-^+NR_{15}R_{16}R_{17}]X^-$ or C_1 - C_5 -fluoroalkyl; R_{10} and R_{11} together are tetra- or pentamethylene either unsubstituted or mono- to trisubstituted by C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, $-SiR_{12}R_{13}R_{14}$, halogen, $-SO_3M$, $-CO_2M$, $-PO_3M$, $-NR_{15}R_{16}$, $[-^+NR_{15}R_{16}R_{17}]X^-$ or C_1 - C_5 -fluoroalkyl, or the group $-PR_{10}R_{11}$ represents a radical of formulae



and R_{12} , R_{13} and R_{14} independently of one another, are C_1 - C_{12} -alkyl or phenyl

R_{15} and R_{16} , independently of one another, are H, C_1 - C_{12} -alkyl or phenyl, or R_{15} and R_{16} together are tetramethylene, pentamethylene or 3-oxa-1,5-pentylene;

R_{17} is H or C_1 - C_4 -alkyl;

M is H or an alkali metal;

X is the anion of a monobasic acid;

halogen is fluorine, chlorine, bromine or iodine.

[0018] The alkyl and alkoxy substituents in question may be, for example, methyl, ethyl, n- and isopropyl, n-, iso- and tert.-butyl, methoxy, ethoxy, n- and isopropoxy, n-, iso- and tert.-butoxy.

[0019] R_{10} and R_{11} in the definition of alkyl may be linear or branched and they preferably contain 1 to 8, most preferably 1 to 4 carbon atoms. Examples of this alkyl are methyl, ethyl, n- and isopropyl, n-, iso- and tert.-butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl and dodecyl. Preference is given to methyl, ethyl, n- and isopropyl, n-, iso- and tert.-butyl. If R_{10} and R_{11} are identical, then as alkyl they most preferably signify isopropyl or tert.-butyl.

[0020] R_{10} and R_{11} , in the definition of cycloalkyl preferably contain 5 to 8, most preferably 5 or 6 ring carbon atoms. Examples of cycloalkyl are cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl and cyclododecyl. Preference is given to cyclopentyl and cyclohexyl, and especially cyclohexyl.

[0021] The cycloalkyl may be substituted, for example by 1 to 3 alkyl or alkoxy substituents; examples of such substituents have already been given. Preference is given to methyl and ethyl, as well as methoxy and ethoxy. Examples of substituted cycloalkyl are methyl- and methoxycyclopentyl and -cyclohexyl.

[0022] R_{10} and R_{11} in the definition of substituted phenyl preferably contain 1 or 2 substituents. Where phenyl contains 2 or 3 substituents, these may be identical or different.

[0023] Examples of alkyl and alkoxy substituents have already been given; preferred alkyl and alkoxy substituents for phenyl are methyl, ethyl and methoxy and ethoxy.

[0024] If the phenyl substituent is halogen, this is preferably -F, -Cl and -Br.

[0025] If the phenyl substituent is C_1 - C_5 -fluoroalkyl, this is partly or wholly fluorinated C_1 - C -alkyl. Examples thereof are the position isomers of mono- to decafluoropentyl, mono- to octafluorobutyl, mono- to hexafluoropropyl, mono-

tetrafluoroethyl and mono- and difluoromethyl. Of the partly fluorinated alkyl radicals, those of formulae $-CF_2H$ and $-CF_2(C_1-C_4\text{-alkyl})$ are preferred in particular. A perfluorinated alkyl is especially preferred. Examples thereof are perfluoropentyl, perfluorobutyl, perfluoropropyl, perfluoroethyl and in particular trifluoromethyl. The fluorine-substituted alkyl groups are preferably bonded in positions 3, 4 and 5.

[0026] R_{12} , R_{13} und R_{14} may be linear or branched alkyl, which preferably contains 1 to 8, most preferably 1 to 4 carbon atoms. Examples of alkyl have already been given. The preferred alkyl is methyl, ethyl, n-propyl, n-butyl and tert.-butyl. The substituent $-SiR_{12}R_{13}R_{14}$ is most preferably trimethylsilyl.

[0027] Of the acidic phenyl substituents $-SO_3M$, $-CO_2M$ and $-PO_3M$, the group $-SO_3M$ and $-CO_2M$ is preferred. M is preferably H, Li, Na and K.

[0028] R_{16} and R_{17} as alkyl preferably contain 1 to 6, most preferably 1 to 4 carbon atoms. The alkyl is preferably linear. Preferred examples are methyl, ethyl, n-propyl and n-butyl. R_{18} as alkyl is preferably methyl.

[0029] X^- as the anion of a monobasic acid is preferably Cl^- , Br^- or the anion of a carboxylic acid or sulphonic acid, for example formate, acetate, trichloroacetate or trifluoroacetate.

[0030] Preferred examples of R_{10} and R_{11} as substituted phenyl are 2-methyl-, 3-methyl-, 4-methyl-, 2- or 4-ethyl-, 2- or 4-isopropyl-, 2- or 4-tert.-butyl-, 2-methoxy-, 3-methoxy-, 4-methoxy-, 2- or 4-ethoxy-, 4-trimethylsilyl-, 2- or 4-fluoro-, 2,4-difluoro-, 2- or 4-chloro-, 2,4-dichloro-, 2,4-dimethyl-, 3,5-dimethyl-, 2-methoxy-4-methyl-, 3,5-dimethyl-4-methoxy-, 3,5-dimethyl-4-(dimethylamino)-, 2- or 4-amino-, 2- or 4-methylamino-, 2- or 4-(dimethylamino)-, 2- or 4- SO_3H -, 2- or 4- SO_3Na -, 2- or 4- $[+NH_3Cl]^-$ -, 3,4,5-trimethylphen-1-yl, 2,4,6-trimethylphen-1-yl, 4-trifluoromethylphenyl or 3,5-di-(trifluoromethyl)phenyl.

[0031] Especially preferred examples of R_{10} and R_{11} are cyclohexyl, tert.-butyl, phenyl, 2- or 4-methylphen-1-yl, 2- or 4-methoxyphen-1-yl, 2- or 4-(dimethylamino)phen-1-yl, 3,5-dimethyl-4-(dimethylamino)phen-1-yl and 3,5-dimethyl-4-methoxyphen-1-yl, but most preferably cyclohexyl, phenyl, 4-methylphen-1-yl and tert.-butyl.

[0032] Another preferred group of compounds is obtained if R_{10} and R_{11} signify unsubstituted phenyl or mono- or disubstituted phenyl.

[0033] A further group of especially preferred compounds is obtained if R_{10} and R_{11} are identical and denote phenyl, cyclohexyl, 2- or 4-methylphen-1-yl, 2- or 4-methoxyphen-1-yl, 2- or 4-(dimethylamino)phen-1-yl, 3,5-dimethyl-4-(dimethylamino)phen-1-yl and 3,5-dimethyl-4-methoxyphen-1-yl; R_{10} and R_{11} are most preferably identical radicals and signify cyclohexyl or phenyl.

[0034] In the context of this invention, functional group means that this group forms a chemical bond by addition or substitution with other functional groups.

[0035] At the functional groups, chain lengthening may be undertaken for example and/or a polymerisable group may be bonded by known methods. Known methods are for example etherification, esterification, amidation, urea formation and urethane formation.

[0036] Processes for the derivatisation of functional groups are known from organic chemistry textbooks (E. Breitmaier, Günther Jung; Organische Chemie II (1983); Georg Thieme Verlag Stuttgart, New York pp.342, 409ff).

[0037] Examples of C-bonded functional groups are the carboxylic acid, carboxylate, carboxylic acid ester, carboxylic acid amide, carboxylic acid halide, cyano, imino, aldehyde, ketone, amine, alcohol, isocyanate, halogen or glycidyl group.

[0038] The functional group may also be a polymerisable group, and in this case is preferably a vinyl group that is unsubstituted or substituted by C_1-C_4 -alkyl.

It may be bonded, for example, by an amide or ester group to the bridging group.

[0039] The polymerisable group may be derived from ethylenically unsaturated alcohols, amines and isocyanates, for example allyl alcohol, allyl amine, allyl isocyanate, croton alcohol; monoesters or monoamides of dicarboxylic acids and unsaturated alcohols and amines; functional styrenes, for example chloromethylstyrene, hydroxystyrene, hydroxyethoxystyrene, styreneamine, styrene-hydroxyethylamine, styrenecarboxylic acid, styrenesulphonic acid, vinyl hydroxyethylether, acrylic acid, methacrylic acid, acrylic and methacrylic acid amide, acrylic- and methacrylic acid- C_2-C_6 -hydroxyalkyl-amide, acrylic- and methacrylic acid- C_2-C_6 -hydroxyalkyl-ester.

[0040] The functional group which is bonded by a bridging group B to one of its carbon atoms preferably signifies an amine, alcohol, isocyanate, carboxylic acid, carboxylic acid ester, carboxylic acid amide, carboxylic acid halide group or a polymerisable group.

[0041] Linkage by means of these functional groups may be carried out by generally known processes. It is fundamentally also possible to transform existing functional groups into other functional groups, for example $-CH_2OH$ groups by oxidation into carboxylic acids, carboxylic acids into amides or halides, amine groups into isocyanate groups, alcohols or amines into carbonates or urethanes. Furthermore, it is possible to react alcohols or amines first of all with halocarboxylic acids (for example chloroacetic acid). Chain extenders, for example epoxides, aziridines, diols, diamines, dicarboxylic acids or dicarboxylic acid esters and diisocyanates, may also be used once or repeatedly in series, thus specifically determining the length of the extending group. These linking methods and processes are known and are described in specialist literature.

[0042] If the functional group FU signifies (O)C-H, (O)C-(C₁-C₁₂)-alkyl, COOH, COCl or COO(C₁-C₆)-alkyl, these groups can also be converted into other functional groups by reduction, transesterification or other known standard reactions from organic chemistry. For example, the aldehyde group is easily converted into an amine group by means of a reaction with an amine and subsequent hydrogenation. Reduction to the alcohol with, for example, LiAlH₄ is likewise possible.

[0043] If the functional group signifies OH, NH₂ or NH(C₁-C₁₂-alkyl), it can be functionalised to an oxyalkylsilyl group by means of a reaction with a compound of formula (R₁₈)_n(R₁₉O)_{3-n}-Si-R₂₀-NCO, whereby R₂₀ is C₁-C₁₂-alkylene, R₁₉ is C₁-C₁₂-alkyl, R₁₈ is C₁-C₁₂-alkyl or OR₁₉ and n is 0, 1 or 2.

[0044] The bridging group B may contain 1 to 30 atoms, preferably 1 to 20 atoms, and most preferably 1 to 12 atoms in the chain, selected from the group C, O, S and N. The bridging group in question is preferably hydrocarbon radicals, that may be interrupted by one or more hetero atoms from the group O, S and N and/or the group C(O).

[0045] The bridging group B may correspond to formula (IV)



wherein X₁ is a direct bond, or X₁ is selected from the group -C(O)-, -O-C(O)-, -SO₂-, -O-SO₂-, -NR₁₀₁-C(O)-, -NR₁₀₁-C(O)-, -NR₁₀₁SO₂- or -NR₁₀₁-SO₂-, wherein

R₁₀₁ is H or C₁-C₃₀-alkyl, C₅- or C₆-cycloalkyl, C₅- or C₆-cycloalkylmethyl or -ethyl, phenyl, benzyl or 1-phenyleth-2-yl and R₁₀₀ is a bivalent bridging group.

[0046] R₁₀₁ defined as alkyl preferably contains 1 to 6, most preferably 1 to 4 carbon atoms. Some examples are methyl, ethyl, n- or isopropyl, butyl, hexyl and octyl. R₁₀₁ defined as cycloalkyl is preferably cyclohexyl, and defined as cycloalkylmethyl is cyclohexylmethyl. In a preferred embodiment, R₁₀₁ is H or C₁-C₄-alkyl.

[0047] The bivalent bridging group R₁₀₀ is preferably a hydrocarbon radical, which preferably contains 1 to 30, more preferably 1 to 18, most preferably 1 to 12, particularly preferably 1 to 8 carbon atoms, and is unsubstituted or mono- or polysubstituted by C₁-C₄-alkyl, C₁-C₄-alkoxy or =O. The hydrocarbon radical may also be interrupted once or many times by hetero atoms selected from the group -O-, -S- and -NR₁₀₁-, whereby R₁₀₁ is preferably H or C₁-C₄-alkyl.

[0048] The bivalent bridging groups R₁₀₀ may be for example C₁-C₂₀-, preferably C₂-C₁₂-alkyls, which can be linear or branched. Some examples are methylene, ethylene, 1,2- or 1,3-propylene, 1,2-, 1,3- or 1,4-butylene, pentylene, hexylene, octylene, dodecylene, tetradecylene, hexadecylene and octadecylene.

[0049] The bivalent bridging group R₁₀₀ may be for example polyoxaalkylene with 2 to 12, preferably 2 to 6, most preferably 2 to 4 oxyalkylene units and 2 to 4, preferably 2 or 3 carbon atoms in the alkylene. It is most preferably polyoxyethylene and polyoxypropylene with 2 to 6 oxyalkylene units.

[0050] The bivalent bridging group R₁₀₀ may be for example C₅-C₁₂-, preferably C₅-C₈-, most preferably C₅- or C₆-cycloalkyl, for example cyclopentylene, cyclohexylene, cyclooctylene or cyclododecylene.

[0051] The bivalent bridging group R₁₀₀ may be for example C₅-C₁₂-, preferably C₅-C₈-, most preferably C₅- or C₆-cycloalkyl-C₁-C₁₂- and preferably -C₁-C₄-alkyl. Some examples are cyclopentyl-C_nH_{2n}- and cyclohexyl-C_nH_{2n}-, wherein n is a number from 1 to 4. -Cyclohexyl-CH₂- is preferred in particular.

[0052] The bivalent bridging group R₁₀₀ may be for example C₅-C₁₂-, preferably C₅-C₈-, most preferably C₅- or C₆-cycloalkyl-(C₁-C₁₂-alkyl)₂- and preferably -(C₁-C₄-alkyl)₂. Some examples are cyclopentyl-(C_nH_{2n})₂ and cyclohexyl-(C_nH_{2n})₂, wherein n is a number from 1 to 4. -CH₂-cyclohexyl-CH₂- is preferred in particular.

[0053] The bivalent bridging group R₁₀₀ may be for example C₆-C₁₄-arylene and preferably C₆-C₁₀-arylene, for example naphthylene or more preferably phenylene.

The bivalent bridging group R₁₀₀ may be for example C₇-C₂₀-aralkylene and preferably C₇-C₁₂-aralkylene. More preferred is arylene-C_nH_{2n}-, wherein arylene is naphthylene and especially phenylene and n is a number from 1 to 4. Examples are benzylene and phenylethylene.

[0054] The bivalent bridging group R₁₀₀ may be for example arylene-(C_nH_{2n})₂-, wherein arylene is preferably naphthylene and especially phenylene, and n is a number from 1 to 4. Examples are xylylene and phenylene(CH₂CH₂)₂-.

[0055] A preferred group of compounds is formed if B signifies unsubstituted linear or branched C₁-C₁₂-alkylene, C₂-C₁₂-alkenylene, C₂-C₁₂-alkynylene, C₅-C₁₂-cycloalkylene, C₅-C₁₂-cycloalkenylene, phenylene, phenylene-(C₁-C₁₂)-alkylene, or B signifies linear or branched C₁-C₁₂-alkylene, C₂-C₁₂-alkenylene, C₂-C₁₂-alkynylene, C₅-C₁₂-cycloalkylene, C₅-C₁₂-cycloalkenylene, phenylene or phenylene-(C₁-C₁₂)-alkylene substituted by C₁-C₄-alkyl, C₁-C₄-alkoxy, halogen or hydroxy, and

FU is halogen, OH, NH₂, NH(C₁-C₁₂)-alkyl, (O)C-H, (O)C-(C₁-C₁₂)-alkyl, COOH, COCl, COO(C₁-C₆)-alkyl, -NCO or a group OC(O)-CR_c=CR_dR_e or OC(NR_f)-CR_c=CR_dR_e, wherein R_c, R_d, R_e and R_f independently of one another, signify hydrogen, C₁-C₆-alkyl or phenyl.

[0056] If group B signifies linear or branched C₁-C₁₂-alkylene, C₂-C₁₂-alkenylene, C₂-C₁₂-alkynylene, C₅-C₁₂-cy-

cloalkylene, C₅-C₁₂-cycloalkenylene, phenylene or phenylene-(C₁-C₁₂)-alkylene substituted by halogen or hydroxy, then at least two functional centres may be present together with the functional group FU, and these can be used for further reactions or for chain extending.

[0057] B is most preferably unsubstituted or halogen- or OH-substituted linear or branched C₁-C₁₂-alkylene.

[0058] Examples of alkylene are methylene, ethylene, the various position isomers of propylene, butylene, pentylene, hexylene, heptylene, octylene, nonylene, decylene, undecylene, dodecylene. Examples of substituted alkyls are 1- or 2-hydroxypropylene, 1-, 2- or 3-hydroxybutylene, the various position isomers of chloropropylene and chlorobutylene. Examples of alkenylene are propenylene, butenylene, pentenylene or hexenylene.


[0059] B as cycloalkylene preferably contains 5 to 8, most preferably 5 or 6 ring carbon atoms.

[0060] Examples of cycloalkylene are cyclopentylene, cyclohexylene, cycloheptylene, cyclooctylene, cyclodecylene, cycloundecylene and cyclododecylene. Preference is given to cyclopentylene and cyclohexylene, and especially cyclohexylene. Cycloalkylene may be substituted by C₁-C₄-alkyl, C₁-C₄-alkoxy, halogen or hydroxy. Examples of such substituents have already been given. Preferred substituents are halogen, OH, methyl and ethyl, as well as methoxy and ethoxy. Examples of substituted cycloalkylene are hydroxycyclohexylene, methyl- and methoxycyclopentylene and -cyclohexylene.

[0061] Examples of cycloalkenylene are cyclopentenylene, cyclohexenylene, cycloheptenylene, cyclooctenylene, cyclodecenylene, cycloundecenylene and cyclododecenylene. Preference is given to cyclopentenylene and cyclohexenylene, and especially cyclohexenylene.

[0062] B when defined as phenylene or phenylene-(C₁-C₁₂)-alkylene, substituted by C₁-C₄-alkyl, C₁-C₄-alkoxy, halogen or hydroxy, preferably contains 1 or 2 substituents. Where phenylene contains 2 or 3 substituents, these may be identical or different. Examples of alkyl and alkoxy substituents have already been given; preferred alkyl and alkoxy substituents for phenylene are methyl, ethyl and methoxy and ethoxy. If the phenylene substituent is halogen, this is preferably -F, -Cl and -Br. Preferred phenylene-(C₁-C₁₂)-alkylene are phenylenepropylene, phenyleneethylene or the benzylene group.

[0063] Some preferred examples of compounds of formula III are those of formula






wherein R₆ and R₇ denote identical or different secondary phosphine groups,

R₂₁ is H, NH₂(CH₂)₃-, (C₁- or C₂-alkoxy)₃Si(CH₂)₃NHC(O)-,

R₂₂ signifies HOCH₂-, H₂NCH₂-, NC-(CH₂)₂-OCH₂-, H₂N-(CH₂)₃-OCH₂- or H₂N-(CH₂)₃-HNCH₂-,

R₂₃ is HO-(CH₂)₂-O- or HO-(CH₂)₃-O-, H₂N-(CH₂)₂-O- or H₂N-(CH₂)₃-O-, and R₂₄ has the same significance as R₂₃ or is methoxy.

[0064] A further aspect of the invention is metal complexes of formulae V, Va and Vb of d-8 metals with the compounds of formula III,



whereby R_6 , R_7 , R_8 and R_9 have the above-mentioned significances and preferences;

Y denotes two monoolefin ligands or one diene ligand;

Me signifies a d-8 metal selected from the group Ir and Rh;

D is -Cl, -Br or -I; and

E is the anion of an oxyacid or complex acid;

X_2 and X'_2 are identical or different and have the significance of D and E, or X_2 and X'_2 are allyl or 2-methylallyl, or X_2 has the significance of D and E and X'_2 is hydride.

[0065] Metal complexes in which Y is 1,5-hexadiene, 1,5-cyclooctadiene or norbornadiene are preferred. In the metal complexes according to the invention, D is preferably -Cl, -Br or -I. In the preferred metal complexes, E is ClO_4^- , CF_3SO_3^- , CH_3SO_3^- , HSO_4^- , BF_4^- , $\text{B}(\text{Phenyl})_4^-$, PF_6^- , SbCl_6^- , AsF_6^- or SbF_6^- .


[0066] Further ruthenium complexes that may be considered are known in literature and are described for example in US 4,691,037, US 4,739,085, US 4,739,084, EP 269395, EP 271310, EP 271311, EP 307168, EP 366390, EP 470756, JP 08081484, JP 08081485, JP 09294932, EP 831099, EP 826694, EP 841343, J. P. Genêt, Arcos Organics Acta, 1 (1995) 4, N.C. Zanetti et al., Organometallics 15 (1996) 860.

[0067] The metal complexes of formulae V, Va or Vb are produced by methods known in literature.

[0068] The compounds of formulae V, Va and Vb represent catalysts that are already homogeneous and can be used, for example, for hydrogenation of unsaturated organic compounds.

[0069] The metal complexes are preferably used for the asymmetric hydrogenation of prochiral compounds with carbon/carbon or carbon/hetero atom multiple bonds, in particular double bonds. Hydrogenations of this type with soluble homogeneous metal complexes are described, for example, in Pure and Appl. Chem., Vol. 68, No. 1, pp. 131-138 (1996).

[0070] The compounds of formula III may be covalently bonded to inorganic or organic carriers in a simple manner. A further aspect of the invention is inorganic or organic polymeric carriers, to which diphosphines of formula III



(III)


are bonded. These are characterised in that they are bonded by the functional group FU of radicals R_8 , R_9 or R_8 and R_9 to the inorganic or polymeric organic carrier, whereby the radicals R_6 , R_7 , R_8 and R_9 have the above-mentioned significances and preferences.

[0071] The diphosphines of formula III are preferably bonded to the surface of these carriers. This has the advantage that catalytically active groups of corresponding d-8 metal complexes are also accessible and no inclusion occurs. In this way, during hydrogenation, less catalyst-containing carrier can also be used.

[0072] If the compounds of formula III are bonded to inorganic carriers, the functional group FU thereof is advantageously first of all reacted with an alkoxysilylalkyl isocyanate, for example a compound of formula (VI)




wherein R_{27} is C_1 - C_{12} -alkylene, R_{26} is C_1 - C_{12} alkyl, R_{25} is C_1 - C_4 -alkyl or OR_{26} and n is 0, 1 or 2; FU in formula III in this case is defined as OH, NH_2 or $NH-(C_1-C_{12})$ -alkyl. Compounds of formula (IIIa) are obtained,




(IIIa)

wherein the group B-FU in radicals R_8 and R_9 is a radical of formula



wherein X_3 signifies -O-, -NH- or $-N(C_1-C_4\text{-alkyl})$, and n , R_{25} , R_{26} and R_{27} have the above-mentioned significances. These compounds are intermediates in the preparation of diphenyldiphosphines bonded to inorganic carriers.

[0073] A further aspect of the invention is a solid inorganic carrier, which is characterised in that it has diphosphine ligands of formula IIIa bonded at the surface by one or two silyl groups of the radical of formula



[0074] During this bonding, 1, 2 or 3 alkoxy groups in the silyl radical can be replaced by bonds.


[0075] The solid carrier in question may be silicates and semi-metal or metal oxides, as well as glass, which preferably exists as powders with average particle sizes of 10 nm to 2000 μm , preferably 10 nm to 1000 μm , most preferably 10 nm to 500 μm . The particles may be both compact and porous. Porous particles preferably have high internal areas, for example 1 to 1200 m^2 , preferably 30 to 600 m^2 . Examples of oxides and silicates are SiO_2 , TiO_2 , ZrO_2 , MgO , NiO , WO_3 , Al_2O_3 , La_2O_3 , silica gels, clays and zeolites. Preferred carriers are silica gels, aluminium oxide, titanium oxide

or glass and mixtures thereof. One example of glass as a carrier is "Controlled Pore Glass", which is available commercially.

[0076] Preparation of the diphosphine ligands of formula IIIa, which are bonded to inorganic carriers, is described in WO 98/01457.

[0077] Owing to the presence of alkoxy silane groups, the compounds of formula IIIa may also be reacted directly to polysiloxanes in a sol-gel process. Reactions of this type have been described for example by U. Deschler et al. in Angew. Chem. 98, (1986), 237-253.

[0078] A further aspect of the invention is organic polymeric carriers, to which diphenylphosphines of formula III



(III)

are bonded by at least one -OH, NH₂ group or functional group FU, whereby the radicals R₆, R₇, R₈ and R₉ have the above-mentioned significances and preferences. These carriers include both polymers which contain as structural elements the diphenyldiphosphines of formula III, which are bonded by at least one -OH, NH₂ group or functional group FU, and polymer particles in which the diphenyldiphosphines of formula III, which are bonded by at least one -OH, NH₂ group or functional group FU, are bonded to functional groups at the surface of the particles.

[0079] The organic polymeric carriers may be uncrosslinked thermoplastic, crosslinked or structurally crosslinked polymers, which contain functional groups.

[0080] The polymers containing functional groups may be either polymers of olefinically unsaturated monomers, for example polyolefins, polyacrylates, polyisoprene, polybutadiene, polystyrene, polyphenylene, polyvinylchloride, polyvinylidene chloride or polyallyl compounds, polyaddition compounds, for example polyurethanes or polyethers, or polycondensated products, for example polyesters or polyamides.

[0081] The monomers which form the polymer are preferably selected from the group styrene, p-methylstyrene or α -methylstyrene, which contain functional groups. Another preferred group of polymers is formed by monomers that are derived from α,β -unsaturated acids, their esters or amides. Particularly preferred are monomers from the group of acrylates and their C₁-C₄-alkylesters, methacrylates and their C₁-C₄-alkylesters, acrylamide and acrylonitrile. An equally preferred group is derived from monomers from the group of acrylates and their C₁-C₄-alkylesters, methacrylates and their C₁-C₄-alkylesters, which contain as structural elements, in bonded form, a hydroxyl group or a primary or secondary amine group as functional groups in the ester group.

[0082] Bonding of the diphenyldiphosphines of formula III to the polymeric carriers may take place in various ways.

[0083] One preferred group of polymerically bonded compounds of formula III is formed in such a way that FU illustrates an olefinically unsaturated radical which is bonded by an ester group OC(O)-CR_c=CR_dR_e or an amide group OC(NR_f)-CR_c=CR_dR_e to the bridging group B, wherein R_c, R_d, R_e and R_f independently of one another, are hydrogen or C₁-C₆-alkyl, and these are used as comonomers in the radical polymerisation of olefinically unsaturated further monomers. Examples and preferences of further monomers are mentioned above. For the remaining radicals of the compounds of formula III, the above-mentioned significances and preferences apply.

[0084] Radical polymerisation is effected in known manner, and a copolymer is obtained which contains diphenyldiphosphine ligands in bonded form.

[0085] Another possibility is a polymer-analogous reaction, such as that described by R. Cullen et. al. in J. of Organometallic Chemistry, 333 (1987), 269-280.

[0086] Polymer-analogous reactions are possible with polycondensates, such as polyesters, polyamines, which contain directly in a side chain or in the polymer chain a further functional group that is capable of condensation. Examples are hydroxyl-group-containing polyesters or polyethers, which can be reacted with compounds of formula III, whereby in this case the functional group FU preferably signifies -COO(C₁-C₁₂)-alkyl or -COCl.

[0087] A further group of preferred polymers that are suitable for polymer-analogous reactions is formed by monomers which contain vinyl alcohol as a homopolymer or vinyl alcohol as a copolymer with vinyl acetate, stearate, benzoate, maleate, vinyl butyral, allyl phthalate, allyl melamine.

[0088] Suitable polymers which are equally preferred for polymer-analogous reactions are formed from phenol and

a C₁-C₄-aldehyde, most preferably from phenol and formaldehyde. The polymers are known as phenol-formaldehyde resins, especially as novolaks, and are available commercially.

[0089] Another preferred group of polymers which are suitable for polymer-analogous reactions is derived from bisglycidyl ethers and diols. These are hydroxyl-functional polyethers, which are produced for example from bisglycidyl ethers and bisphenol A. These polyepoxides may be built up from diepoxide comonomers with preferably 6 to 40, most preferably 8 to 30 carbon atoms, and diols as comonomers with preferably 2 to 200, most preferably 2 to 50 carbon atoms. One preferred group derived therefrom is formed from monomers, which build up a polymer from cyclic C₃-C₆-ethers or C₂-C₆-alkylene glycols with bisglycidyl ethers. The bisglycidyl ethers may be aromatic, aliphatic or cycloaliphatic.

[0090] Further preferred polymers with hydroxyl groups as functional groups are polysaccharides. Especially preferred are partial cellulose acetates, propionates or butyrates, partial cellulose ethers, starch, chitin and chitosan.

[0091] Further polymers are derived from polymers having reducible groups, for example nitrile groups, ketone groups, carboxylic acid esters and carboxylic acid amides.

[0092] Insoluble polymers may also be used in the reaction medium, and these are functionalised at the surface with hydroxyl or amine groups by means of a chemical or physical process. For example, partially unsaturated polymers are provided at their surface with hydroxyl groups by means of oxidation, e.g. with hydrogen peroxide. Another possibility is plasma treatment in, for example, an oxygen atmosphere, nitrogen atmosphere or ammonia atmosphere. The polymers are preferably present as powders. Of these carriers, polystyrene is preferred in particular, and this is subsequently functionalised by known methods with hydroxyl, amino or hydroxymethyl groups.

[0093] An especially preferred group is formed by a polymeric organic material with structural elements, in which at least one isocyanate group FU in compounds of formula III is bonded to hydroxyl or amine groups, forming a urethane or urea bond, whereby the hydroxyl or amine groups are bonded directly or in a side chain of the polymer chain. Monomers of formula III with isocyanate groups are obtainable in a simple manner by reacting diisocyanates with amine- or hydroxyl-functional compounds of formula III.

[0094] The diphenyldiphosphine radicals of formula III may be present as enantiomer mixtures. The radicals are preferably present in the form of the optically active isomers.

[0095] One preferred group of immobilised polymers according to the invention is that in which hydroxy- or amino-functional polymers are first of all reacted with one isocyanate group of diisocyanates and then the second isocyanate group is reacted with a hydroxy- or amino-functional diphosphine of formula III.

[0096] The choice of diisocyanate is in itself not critical. Suitable diisocyanates that are available on a large scale are described for example in Houben Weyl, Makromolekulare Stoffe, volume E 20, pages 1587 to 1583, 1987 edition.

[0097] Preference is given to diisocyanates with a bridging group Q, which is selected from the group linear or branched, aliphatic C₂-C₂₀-alkyl which is unsubstituted or mono- to polysubstituted by C₁-C₆-alkyl, C₁-C₆-alkoxy; C₃-C₈-cycloalkyl or heterocycloalkyl which is unsubstituted or mono- to polysubstituted by C₁-C₆-alkyl, C₁-C₆-alkoxy; linear or branched aliphatic C₂-C₂₀-alkyl unsubstituted or substituted by C₁-C₆-alkyl, C₁-C₆-alkoxy and interrupted by C₃-C₈-cycloalkyl or heterocycloalkyl which is unsubstituted or substituted by C₁-C₆-alkyl, C₁-C₆-alkoxy; phenyl, naphthyl, biphenyl or C₃-C₁₀-heteroaryl either unsubstituted or mono- to polysubstituted by C₁-C₆-alkyl, C₁-C₆-alkoxy; linear or branched aliphatic C₂-C₂₀-alkyl which is unsubstituted or substituted by C₁-C₆-alkyl, C₁-C₆-alkoxy and is interrupted by phenyl, naphthyl or C₃-C₁₀-heteroaryl.

[0098] Heterocycloalkyl is e.g. pyrrolidine, piperidine, morpholine, oxazolidine, dioxolan or an isocyanuric acid triester group.

[0099] Heteroaryl is for example pyridine, pyrimidine, pyrrole, furan, imidazole, pyrazole or triazine.

[0100] Especially preferred diisocyanates are 1,6-bis-[isocyanate]-hexane, 5-isocyanate-3-(isocyanatemethyl)-1,1,3-trimethylcyclohexane, 1,3-bis-[5-isocyanate-1,3,3-trimethyl-phenyl]-2,4-dioxo-1,3-diazetidene, 3,6-bis-[9-isocyanate-nonyl]-4,5-di-(1-heptenyl)-cyclohexene, bis-[4-isocyanate-cyclohexyl]-methane, traps-1,4-bis-[isocyanate]-cyclohexane, 1,3-bis-[isocyanatemethyl]-benzene, 1,3-bis-[1-isocyanate-1-methyl-ethyl]-benzene, 1,4-bis-[2-cyanate-ethyl]-cyclohexane, 1,3-bis-[isocyanate-methyl]-cyclohexane, 1,4-bis-[1-isocyanate-1-methylethyl]-benzene, bis-[isocyanate]-isododecylbenzene, 1,4-bis-[isocyanate]-benzene, 2,4-bis-[isocyanate]-toluene, 2,6-bis-[isocyanate]-toluene, 2,4-/2,6-bis-[isocyanate]-toluene, 2-ethyl-1,2,3-tris-[3-isocyanate-4-methyl-anilinocarbonyloxy]-propane, N,N'-bis-[3-isocyanate-4-methylphenyl]-urea, 1,4-bis-[3-isocyanate-4-methylphenyl]-2,4-dioxo-1,3-diazetidene, 1,3,5-tris-[3-isocyanate-4-methylphenyl]-2,4,6-trioxohexahydro-1,3,5-triazine, 1,3-bis-[3-isocyanate-4-methylphenyl]-2,4,5-trioximidazolidine, bis-[2-isocyanate-phenyl]-methane, (2-isocyanatephenyl)-(4-isocyanate-phenyl)-methane, bis-[4-isocyanate-phenyl]-methane, 2,4-bis-[4-isocyanate-benzyl]-1-isocyanatbenzene, [4-isocyanate-3-(4-isocyanate-benzyl)-phenyl]-[2-isocyanate-5-(4-isocyanate-benzyl)-phenyl] methane, tris-[4-isocyanate-phenyl]-methane, 1,5-bis-[isocyanate]-naphthalene, or 4,4'-bis[isocyanate]-3,3'-dimethyl-biphenyl.

[0101] Particularly preferred diisocyanates are 1,6-bis[isocyanate]-hexan, 5-isocyanate-3-(isocyanate-methyl)-1,1,3-trimethylcyclohexane, 2,4-bis-[isocyanate]-toluene, 2,6-bis-[isocyanate]-toluene, 2,4-/2,6-bis-[isocyanate]-toluene or bis-[4-isocyanate-phenyl]-methane.

[0102] The polymers to be used according to the invention, which contain hydroxyl groups and amine groups, may be uncrosslinked thermoplastic, crosslinked or structurally crosslinked polymers. Examples of hydroxyl-group-containing polymers are those mentioned above.

[0103] The polymers to be used according to the invention are known *per se*, partly commercial or may be produced by known polymerisation processes or by subsequent modification of polymers.

[0104] The polymeric organic materials preferably have a molecular weight of 5000 to 5,000,000 daltons, most preferably 50,000 to 1,000,000 daltons.

[0105] A preferred sub-group of polymeric organic materials is highly crosslinked macroporous polystyrene or polyacrylate.




[0106] Another preferred group of polymers is formed by weakly crosslinked polystyrene. An example thereof is polystyrene crosslinked with 1-5% divinylbenzene.

[0107] The particle size of the polymeric organic materials is preferably 10 μm to 2000 μm .

[0108] The highly crosslinked polymeric organic materials preferably have a specific area of 20 m^2/g to 1000 m^2/g , most preferably 50 m^2/g to 500 m^2/g , determined by the BET method.

[0109] Production of the diphenyldiphosphines that are bonded to inorganic or organic carriers may be effected analogously to the processes described in WO 98/01457.

[0110] A further aspect of the invention is the d-8 metal complexes of inorganic or organic polymeric carriers, to which diphenyldiphosphines of formula VII, VIIa or VIIb



are bonded by at least one HO-, H₂N- group or functional group FU, whereby the radicals R₆, R₇, R₈, R₉, Me, E, Y, X₂ and X₂' and the carrier have the above-mentioned significances and preferences.

[0111] Production of metal complexes with the polymeric carrier may take place by methods known in literature for the production of analogous homogeneous catalysts.

[0112] A further aspect of the invention is diphenyldiphosphine ligands of formula III and their d-8 metal complexes of formulae VII, VIIa and VIIb with a molecular weight of preferably less than 5000 daltons, which contain solubility-enhancing or adsorption-facilitating groups bonded to at least one HO-, H₂N- or functional group FU, which groups can be separated by extraction with immiscible liquids or by adsorption on a carrier. During extraction, it is preferable to use two immiscible liquids. When using such diphenyldiphosphines and the d-8 metal complexes thereof, there is almost no loss of metal or ligand. Therefore, using these extractable or adsorbable catalysts, large-scale hydrogenation may be carried out especially economically. A preferred group of soluble compounds is those that are soluble in aqueous media.

[0113] Suitable immiscible liquids that may be mentioned are, for example, water and organic solvents that are immiscible with water, such as alkanes (for example hexane), chlorinated alkanes (for example methylene chloride, chloroform), aryls (for example toluene, benzene, xylene) or esters (for example ethyl acetate) or organic solvent systems such as fluorinated hydrocarbons and hydrocarbons.

[0114] Carriers that are suitable for adsorption are metal oxides, for example silica gel, aluminium oxide, or reversed phase silica gel, polar and apolar polymers and ion exchangers (preferably for ligands with charged radicals).

[0115] Suitable solubility-enhancing or adsorption-facilitating radicals may be taken from the publication I.T. Horváth et al. in Science, Vol. 266, pages 72-75 (1994).

[0116] Preferred solubility-enhancing radicals for extractable diphenyldiphosphines are, for example, lipophilic radicals which are derived from alkanes having a molecular weight of between 100 and 2000 daltons, or also hydrophilic, optionally charged radicals, which are derived from sugars, or from polymers, for example polyvinyl alcohols, polyacrylic acids, polyethylene glycols, polyvinyl toluene or dendrimers.

[0117] Preferred adsorption-facilitating radicals are, for example, lipophilic radicals which are derived from alkanes having a molecular weight of between 100 and 2000 daltons, and also fluoroalkanes.

[0118] Further examples of extractable or adsorbable radicals are those that are derived from polyethylene glycols, polyhydroxy hydrocarbons, polyamino hydrocarbons and the ammonium salts thereof, polycarboxyl hydrocarbons and the alkali metal salts thereof, polyhydroxy hydrocarbons or polyamino hydrocarbons, which are reacted with halocarboxylic acids, polyvinyl alcohols, polyaryl acids and the alkali metal salts thereof, higher alkanes and perfluoroalkanes.

[0119] The metal complexes according to the invention are eminently suitable as catalysts for the hydrogenation of organic double and triple bonds. Examples are compounds which contain the groups C=C, C=N, C=O, C=C-N or C=C-O (see for example K. E. König, The Applicability of Asymmetric Homogeneous Catalysis, in James D. Morrison (ed.), Asymmetric Synthesis, Vol. 5, Academic Press, 1985). The metal complexes according to the invention are especially suitable for the enantioselective hydrogenation of compounds having prochiral carbon double bonds and carbon/hetero atom double bonds. Examples of such compounds are prochiral alkenes, imines and ketones.

[0120] After the reaction, the catalysts according to the invention may be practically completely separated from the reaction mixture in a simple manner, for example by decanting, centrifuging, filtration, ultrafiltration, extraction or adsorption, and reused. One particular advantage of this is that they can be reused several times without any notable losses of activity or selectivity. The catalysts which are functionalised and immobilised according to the invention often have improved optical yields when compared with the previously known diphenylphosphine catalysts.

[0121] A further aspect of the invention is therefore the use of the metal complexes of d-8 metals according to the invention as heterogeneous or homogeneous catalysts for the asymmetric hydrogenation of prochiral compounds with carbon double bonds or carbon/hetero atom double bonds. The metal complexes are preferably used for the asymmetric hydrogenation of prochiral compounds with carbon double bonds or carbon/hetero atom double bonds, especially the Ir complexes for the hydrogenation of prochiral ketimines.

[0122] A further aspect of the invention is a process for the asymmetric hydrogenation of compounds with carbon double bonds or carbon/hetero atom double bonds, which is characterised in that the compounds are reacted at a temperature of -20 to 80 °C and at a hydrogen pressure of 10^5 to 2×10^7 Pa in the presence of catalytic amounts of one or more metal complexes according to the invention.

[0123] Catalysts are preferably employed in amounts of 0.0001 to 10 mol %, more preferably 0.001 to 10 mol %, most preferably 0.01 to 5 mol %, based on the compound to be hydrogenated.

[0124] Hydrogenation may be carried out continuously or intermittently in various types of reactor. Preference is given to the reactors which allow comparatively favourable blending and good heat removal, e.g. loop reactors. This type of reactor has proved to be particularly effective when using small amounts of catalyst.

[0125] The hydrogenated organic compounds that may be produced according to the invention are active substances or intermediates for producing such substances, especially in the field of pharmaceutical and agrochemical production. For example, o,o-dialkylarylketamine derivatives, especially those with alkyl and/or alkoxyalkyl groups, have fungicidal activity, especially herbicidal activity. The derivatives in question may be amine salts, acid amides, e.g. of chloroacetic acid, tertiary amines and ammonium salts (see e.g. EPA-0 077 755 and EP-A-0 115 470).

[0126] The following examples illustrate the invention.

A) Preparation of halomethyldiphenyl diiodides

Example A1: (R)-[6-bromomethyl-6'-methyldiphen-2,2'-diyl]bisiodide and (R)-[6,6'-dibromomethyldiphen-2,2'-diyl]bisiodide

[0127] 3 g of (R)-[6,6'-dimethyldiphen-2,2'-diyl]bisiodide (6.91 mmols), 1.23 g of N-bromosuccinimide (6.91 mmols) and 30 mg of α,α' -azoisobutyronitrile are mixed under argon with 45 ml of CCl_4 and heated under reflux for one hour. After adding a spatula tip of α,α' -azobutyronitrile, heating under reflux takes place again for 24 hours. A spatula tip of α,α' -azobutyronitrile is again added and heating under reflux takes place for a further 14 hours. The mixture is concentrated on a rotary evaporator and extracted with $\text{H}_2\text{O}/\text{CH}_2\text{Cl}_2$. The organic phase is separated, dried (MgSO_4), and concentrated on a rotary evaporator. After chromatography on silica gel (hexane \rightarrow hexane/ethyl acetate (ea) = 5/1),

1.98 g of product is isolated as an oil (56%, 3rd fraction).

$[\alpha]_D = +63$ (CH_2Cl_2 , $c = 0.82$).

$^1\text{H-NMR}$: 7.96 - 6.68 (*m*, 6 arom. H); 4.31 (*d*, $J = 10.5$; 1 H CH_2Br); 4.08 (*d*, $J = 10.5$; 1 H CH_2Br); 2.17 (*s*, 1 CH_3).

[0128] (R)-[6,6'-dibromomethyldiphen-2,2'-diyl]bisiodide is obtained as a by-product in a yield of 9%. $[\alpha]_D + 52.3$ (CHCl_3 , $c = 1.2$) $^1\text{H-NMR}$: 7.97 - 7.2 (*m*, 6 arom. H); 4.26 (*d*, $J = 10.9$; 2 CHBr); 4.19 (*d*, $J = 10.9$, 2 CHBr).

B) Preparation of hydroxymethyldiphenyl bisiodides


Example B1: Preparation of (R)-[6-hydroxymethyl-6'-methyl-diphen-2,2'-diyl]-bisiodide

[0129] 14.38 g of the compound according to example A1 (0.028 mols) are dissolved in 350 ml of dioxane and mixed with a KOH solution (18.6 g in 300 ml of H_2O). The mixture is heated over night under reflux, whereby a clear solution is obtained. After cooling to room temperature, the mixture is extracted with CH_2Cl_2 , the organic phase separated, dried (MgSO_4) and concentrated on a rotary evaporator. After recrystallisation in cyclohexane, 12.1 g of white crystals are obtained (95.6%). Melting point: 120 °C; $[\alpha]_D = -25.3$ (CHCl_3 , $c = 0.73$).

$^1\text{H-NMR}$: 7.96 - 7.02 (*m*, 6 arom. H); 4.35 (*d*, $J = 13.8$; 1 H CH_2O); 4.24 (*d*, $J = 13.8$; 1 H CH_2O); 2.03 (*s*, 1 CH_3); 1.72 (*s*, 1 HO).

C) Preparation of functional diphenyldiphosphines

[0130] Example C1: Preparation of




0.47 g of Cs_2CO_3 (1.4 mmols) are mixed with 60 ml of absolute CH_3CN and heated under reflux. 100 mg of (R)-6,6'-dihydroxy-diphenyl-2,2'-diphenyldiphosphine (0.18 mmols), dissolved in 5 ml of CH_3CN , and 24.7 mg of epibromohydrin (0.18 mmols), dissolved in 5 ml of CH_3CN , are slowly added dropwise over the course of 14 hours using a spray pump. Afterwards, a further 8 mg of epibromohydrin (0.06 mmols) are added and heating continues for 2 hours under reflux. The solid residue is filtered and the solution concentrated on a rotary evaporator. After chromatography on silica gel (hexane/ea = 4/1), 40 mg of a white solid are obtained (36.3 %).

When reacting 100 mg of (R)-6,6'-dihydroxy-diphenyl-2,2'-diphenyldiphosphine (0.18 mmols) with 39 mg of 1,3-dibromo-2-propanol (0.18 mmols) in the same way as before, 40 mg of the same product are isolated. Melting point: 122 °C; $[\alpha]_D = -290$ (CHCl_3 , $c = 1$).

$^1\text{H-NMR}$: 7.6 - 6.85 (*m*, 26 arom. H); 4.48 (*d*, $J = 12.9$; 1 H CH_2O); 4.33 (*dd*; $J = 11.2, 5.6$; 1 H CH_2O); 4.09 (*d*, $J = 11.2$, 1 H CH_2O); 3.98 (*dd*, $J = 12.9, 2.6$; 1 H CH_2O); 3.74 (*m*; 1 HCOH); 2.04 (*d*, $J = 10.3$, 1 HO); $^{31}\text{P}\{^1\text{H}\}\text{-NMR}$: -9.6 (*s*)

[0131] Example C2: Preparation of




[0132] The reaction of 0.4 mg of (S)-6,6'-dihydroxy-diphenyl-2,2'-diphenyldiphosphine (0.72 mmols) with 98 mg of epibromohydrin in analogous manner to that above yields 144 mg of the title compound (32.7 %). Melting point: 122 °C; $[\alpha]_D = +289$ (CHCl_3 , $c = 0.8$).

$^1\text{H-NMR}$ and $^{31}\text{P}\{^1\text{H}\}\text{-NMR}$ identical to that of compound C1.

[0133] Example C3: Preparation of

¹H-NMR and ³¹P{¹H}-NMR identical to that of compound C1.

[0135] Example C4: Preparation of



¹H-NMR (CDCl₃): (2 isomers by reversal of the propyl-triethoxysilane group on nitrogen in a ratio of 6.3 to 1) 7.75-6.65 (*m*, 26 arom. H); 4.85 (*m*, 1 NH, 1 OCH); 4.7 (br. s, 1 NH)_{ki}; 4.45 (br. *dd*, *J* = 11.6, 3.6; 1 H CH₂OPh); 4.3 (br. *d*, *J* = 11.6; 1 H CH₂Ph); 4.1 (br. *dd*, *J* = 11.6, 7.1; 1 H CH₂OPh); 3.95 (*m*, 1 H CH₂OPh); 3.85 (*q*, *J* = 7; 3 CH₂O); 3.15 (*m*, 1 CH₂N)_{gr}; 2.82 (*m*, 1 CH₂N)_{ki}; 1.63 (*m*, 1 CH₂CH₂N)_{gr}; 1.50 (*m*, 1 CH₂CH₂N)_{ki}; 1.25 (*t*, *J* = 7; 3 CH₃); 0.64 (*dd*, *J* = 8.3, 7.2; 1 CH₂Si)_{gr}; 0.55 (*m*, 1 CH₂Si)_{ki}.

 $^{31}\text{P}\{^1\text{H}\}$ -NMR: -10.11 (s)_{gr}; -9.88 (s)_{kl}

Example C5:

¹H-NMR and ³¹P{¹H}-NMR are identical to compound C4.

Example C6: Preparation of (R)-[6-hydroxymethyl-6'-methyl-diphen-2,2'-diyl]-bis-(diphenylphosphine)

a) (R)-[6-(*t*-butyl-dimethylsilyloxomethyl)-6'-methyl-biphenyl-2,2'-diyl]-bisiodide

¹H-NMR: 7.9 - 7.02 (*m*, 6 arom H); 4.35 (*d*, *J* = 14.4; 1 H CH₂O); 4.15 (*d*, *J* = 14.4; 1 H CH₂O); 2.02 (*s*, 1 CH₃); 0.91 (*s*, 3 CH₃); 0.03 (*s*, 1 CH₃Si); 0.01 (*s*, 1 CH₃Si).

b) (R)-[6-(*t*-butyl-dimethylsilyloxomethyl)-6'-methyl-biphenyl-2,2'-diyl]-bis-(diphenylphosphine)

16

orator and extracted with $\text{H}_2\text{O}/\text{CH}_2\text{Cl}_2$. The organic phase is separated, dried (MgSO_4), and concentrated on a rotary evaporator. After chromatography on silica gel (hexane/ea = 80/1), 1.62 g of a white solid are isolated (51.8 %). Melting point: 97°C ; $[\alpha]_D = +7.3$ (CHCl_3 , $c = 0.91$).

^1H -NMR: 7.51 - 7.06 (m, 26 arom. H); 3.75 (d, $J = 13.3$, 1 H CH_2O); 3.45 (d, $J = 13.3$, 1 H CH_2O); 1.45 (s, 1 CH_3Ph); 0.81 (s, 3 CH_3); -0.18 (s, 1 CH_3Si); -0.20 (s, 1 CH_3Si).

$^{31}\text{P}\{^1\text{H}\}$ -NMR: -14.1 (d, $J = 30$); -14.8 (d, $J = 30$).

c) (R)-[6-hydroxymethyl-6'-methyl-diphen-2,2'-diyl]-bis-(diphenylphosphine)

[0139] 1.522 g of the above compound b) (2.4 mmols) are dissolved in 25 ml of absolute tetrahydrofuran (THF) and mixed with 67 ml of tetrabutylammonium fluoride solution (1 M in THF, 6.7 mmols). The solution is stirred over night (16 hours) and then concentrated in a rotary evaporator. After extraction with water/tert.-butylmethylether, the organic phase is separated, dried (MgSO_4) and concentrated on a rotary evaporator. After chromatography on silica gel (hexane/ea = 3/1), 1.11 g of a white solid are obtained, which is recrystallised in methanol. A crystalline product is obtained (white needles); yield 0.91 g (71.8 %). Melting point: 215°C ; $[\alpha]_D = -20.4$ (CHCl_3 , $c = 0.84$).

^1H -NMR: 7.40 - 7.05 (m, 26 arom. H); 3.86 (dd, $J = 13.7$, 9; 1 H CH_2O); 3.44 (dd, $J = 13.7$, 4.3; 1 H CH_2O); 1.35 (s, 1 CH_3); 0.33 (dd, $J = 9$, 4.3; 1 HO).

$^{31}\text{P}\{^1\text{H}\}$ -NMR: -13.6 (d, $J = 40.3$); -13.9 (d, $J = 40.3$).

Example C7: Preparation of (R)-[6-hydroxymethyl-6'-methyl-diphen-2,2'-diyl]-bis-(dicyclohexylphosphine)

a) (R)-[6-(t.-butyl-dimethylsilyloxomethyl)-6'-methyl-biphenyl-2,2'-diyl]-bis-(dicyclohexylphosphine)

[0140] 1.21 g of compound C6(a) (2.14 mmols) are dissolved in 20 ml of absolute diethylether (Et_2O) and cooled to -78°C . 4.5 ml of lithium butyl solution (1.6 M in hexane, 4.5 mmols) are then slowly added dropwise, and the solution is stirred for 2 hours at -78°C . 1.24 g of chlorodicyclohexylphosphine (5.36 mmols) are added dropwise, and the mixture is then heated to room temperature. After 20 hours under reflux, the mixture is extracted with $\text{H}_2\text{O}/\text{CH}_2\text{Cl}_2$. The organic phase is separated, dried (MgSO_4), and concentrated on a rotary evaporator. After chromatography on silica gel (hexane/ea = 100/1), 1.35 g of a colourless oil are isolated (89 %). This oil is used further without further purification. $^{31}\text{P}\{^1\text{H}\}$ -NMR: -8.5 (d, $J = 20.8$); -9.2 (d, $J = 20.8$).

b) (R)-[6-hydroxymethyl-6'-methyl-diphen-2,2'-diyl]-bis-(dicyclohexylphosphine)

[0141] 1.348 g of the above compound a) (1.91 mmols) are dissolved in 25 ml of absolute THF and mixed with 5.7 ml of tetrabutylammonium fluoride solution (1 M in THF, 5.7 mmols). The solution is stirred over night (16 hours) and then concentrated in a rotary evaporator. After extraction with water/tert.-butylmethylether, the organic phase is separated, dried (MgSO_4) and concentrated on a rotary evaporator. After chromatography on silica gel under argon (hexane/ea = 4/1), 0.3 g of a white solid are isolated (26 %). M.p.: 213°C .

^1H -NMR (CD_2Cl_2): 7.56 - 7.21 (m, 6 arom. H); 4.21 (d, $J = 12.9$, 1 H CH_2O); 4.10 (dd, $J = 12.9$, 1; 1 H CH_2O); 2.15-0.80 (m, 45 H: OH+Cy); 1.91 (s, 1 CH_3). $^{31}\text{P}\{^1\text{H}\}$ -NMR: -8.9 (s)

Example C8: Preparation of (R)-[6-aminomethyl-6'-methyl-diphen-2,2'-diyl]-bis-(diphenylphosphine)

a) (R)-[6-phthalimid-N-yl-methyl-6'-methyl-diphen-2,2'-diyl]-bis-(diphenylphosphine)

[0142] 0.173 ml (1 mmol) of azodicarboxylic acid diethylester (purity 90%) are added dropwise to a solution of 380 mg (0.67 mmols) of compound C6(c), 352 mg (1.34 mmols) of triphenylphosphine and 128 mg (0.87 mmols) of phthalimide in 15 ml of THF, and the solution stirred over night. The solution is then concentrated on a rotary evaporator, the residue extracted in water/methylene chloride, the organic phase dried with sodium sulphate and concentrated on a rotary evaporator. After purification by chromatography (silica gel Merck 60; eluant = methylene chloride), 130 mg of product are obtained (white powder, yield 28%).

^1H -NMR (CDCl_3): 7.85 - 7.6 (m, 4H, phthalimide), 7.5 - 7.0 (m, 26 arom. H), 4.0 (d, $J = 14.3$, 1 H, $\text{CH}_2\text{-N}$), 3.8 (d, $J = 14.3$, 1 H, $\text{CH}_2\text{-N}$), 1.59 (s, 3H, CH_3). $^{31}\text{P}\{^1\text{H}\}$ -NMR: -15.4 (s)

b) (R)-[6-aminomethyl-6'-methyl-diphen-2,2'-diyl]-bis-(diphenylphosphine)

[0143] 128 mg (0.184 mmols) of the above compound a) in 10 ml of ethanol are heated under reflux for 4 hours with 0.15 ml of hydrazine hydrate. The suspension which is obtained upon cooling is filtered, the residue washed with ethyl

acetate, and the filtrate concentrated on a rotary evaporator. After purification by chromatography (silica gel Merck 60; eluant = ethanol/triethylamine 250:1), 95 mg of product are obtained (white powder, yield 91%).

$^1\text{H-NMR}$ (CDCl_3): 7.45 - 7.0 (m, 26 arom. H), 3.05 (d, $J = 12.3$, 1 H, $\text{CH}_2\text{-N}$), 2.69 (d, $J = 12.3$, 1 H, $\text{CH}_2\text{-N}$), 1.33 (s, 3H, CH_3). $^{31}\text{P}\{^1\text{H}\}\text{-NMR}$: -14.0 (s)

Example C9: Preparation of (R)-[6- β -cyanoethyloxymethyl-6'-methyl-diphen-2,2'-diyl]-bis-(diphenylphosphine)

[0144] 0.107 ml (1.63 mmols) of acrylonitrile and then 0.005 ml of a 40% aqueous KOH solution are added to a solution of 300 mg (0.53 mmols) of compound C6(c) in 3 ml of dioxane, and the reaction solution is subsequently stirred for 24 hours. Then, the solution is concentrated on a rotary evaporator and the product is purified by chromatography (silica gel Merck 60, eluant = methylene chloride). 313 mg of product are obtained (white powder, yield 95%).

$^1\text{H-NMR}$ (CDCl_3): 7.45 - 7.0 (m, 26 arom. H), 3.68 (d, $J = 12.5$, 1 H, $\text{Ph-CH}_2\text{-O}$), 3.19 (d, $J = 12.5$, 1 H, $\text{Ph-CH}_2\text{-O}$), 3.13 - 2.9 (m, 2H, $\text{O-CH}_2\text{-CH}_2$), 2.3 (t, 2H, $\text{CH}_2\text{-CN}$), 1.38 (s, 3H, CH_3). $^{31}\text{P}\{^1\text{H}\}\text{-NMR}$: -14.7 (s)

Example C10: Preparation of (R)-[6- γ -aminopropoxyloxymethyl-6'-methyl-diphen-2,2'-diyl]-bis-(diphenylphosphine)

[0145] A suspension of 325 mg (0.52 mmols) of compound C9 and 60 mg of LiAlH_4 in 5 ml of diethylether is stirred over night. After adding 20 ml of diethylether, the mixture is hydrolysed with ca. 0.5 ml of water and the whole mixture is subsequently dried with sodium sulphate. The solution is concentrated by evaporation and the residue purified by chromatography (silica gel Merck 60, eluant = ethanol/triethylamine 100:1). 157 mg of product are obtained (yield 45%). Compound C6(c) is formed as a by-product.

$^1\text{H-NMR}$ (CDCl_3): 7.45 - 7.0 (m, 26 arom. H), 3.6 (d, $J = 12.5$, 1 H, $\text{Ph-CH}_2\text{-O}$), 3.17 (d, $J = 12.5$, 1 H, $\text{Ph-CH}_2\text{-O}$), 2.96 (m, 2H, $\text{O-CH}_2\text{-CH}_2$), 2.65 (t, $J = 6.7$, 2H, $\text{CH}_2\text{-N}$), 1.52 (m, 2H, $\text{O-CH}_2\text{-CH}_2$), 1.4 (s, 3H, CH_3).


Example C11: Preparation of (R)-[6- γ -trimethoxysilylpropylaminocarbonyloxymethyl-6'-methy-diphen-2,2'-diyl]-bis-(diphenylphosphine)

[0146] 209 mg of compound C6(c) (0.37 mmols) are dissolved in 5 ml of absolute CH_2Cl_2 . To this solution are added dropwise 322 mg of 3-(isocyanatopropyl)-triethoxysilane (0.49 mmols) and 5 ml of dibutyltin dilaurate. The solution is heated under reflux over night and then concentrated in a rotary evaporator. After chromatography on silica gel (hexane/ea = 4/1), 260 mg of a colourless oil are obtained (87 %). $[\alpha]_D = 21.7$ (CHCl_3 , $c = 1.12$).

$^1\text{H-NMR}$ (CDCl_3 ; 400.16 MHz): (2 isomers by reversal of the propyl-triethoxysilane group on nitrogen in a ratio of 3.6 to 1) 7.40-7.10 (m, 24 arom. H); 7.06 (m, 2 arom. H); 4.93 (br. s, 1 NH_{ki}); 4.63 (br. t, $J = 5.7$; 1 NH_{gr}); 4.25 (d, $J = 12.9$; 1 H CH_2O); 4.13 (q, $J = 7.3$); 3.86 (q, $J = 7$, 3 $\text{CH}_2\text{OSi}_{\text{ki}}$); 3.84 (d, $J = 7$; 1 H CH_2O); 3.83 (d, $J = 7$, 3 $\text{CH}_2\text{OSi}_{\text{gr}}$); 3.21 (q, $J = 6.6$; 1 $\text{CH}_2\text{N}_{\text{ki}}$); 3.09 (q, $J = 6.7$; 3 $\text{CH}_2\text{N}_{\text{gr}}$); 1.66 (m, 1 $\text{CH}_2\text{CH}_2\text{N}_{\text{ki}}$); 1.57 (m, 1 $\text{CH}_2\text{CH}_2\text{N}_{\text{gr}}$); 1.43 (s, 1 CH_3Ph); 1.26 (t, $J = 7$; 3 CH_3_{ki}); 1.24 (t, $J = 7$; 3 CH_3_{gr}); 0.67 (m, 1 $\text{CH}_2\text{Si}_{\text{ki}}$); 0.60 (m, 1 $\text{CH}_2\text{Si}_{\text{gr}}$). $^{31}\text{P}\{^1\text{H}\}\text{-NMR}$: -13.2 (d, $J = 34.7$); -13.9 (d, $J = 34.7$).

Example C12: Preparation of a water-soluble diphosphine of formula

[0147]



a) A solution of 380 mg (0.9 mmols) of tris-1,2,3-(ethoxycarbonylethyloxymethyl)-2-aminopropane, that has been dried by azeotropic distillation in toluene, in 8.5 ml of absolute toluene is added to 146 mg (0.9 mmols) of 1,1'-carbonyl-diimidazole, and the reaction mixture is stirred for 3 days at room temperature. To this mixture are added first of all a solution of 300 mg (0.53 mmols) of compound C6(c) in 8 ml of absolute toluene and then 20 mg of dibutyltin dilaurate, and the reaction mixture is stirred for 2 days at 100°C. The reaction mixture is then concentrated on a rotary evaporator and purified by chromatography (silica gel Merck 60, eluant: hexane/ethyl acetate 2:1) 415

mg of product are obtained (almost solid oil, yield 80%).

$^1\text{H-NMR}$: 7.40 -7.0 (*m*, 26 arom. H); 4.95 (*s*, 1H, NH), 4.2 (*d*, $J = 14$, 1 H, CH_2O); 4.11 (*q*, 6H, CH_2CH_3), 3.77 (*d*, $J = 14$, 1 H CH_2O); 3.63 (*t*, 6H, $\text{O-CH}_2\text{CH}_2$), 3.08 (*s*, 6H, $\text{C-CH}_2\text{-O}$), 2.5 (*t*, 6H, $\text{O-CH}_2\text{CH}_2$), 1.38 (*s*, 3 H, Ph-CH_3), 1.22 (*t*, 9H, CH_2CH_3).

$^{31}\text{P}\{^1\text{H}\}$ -NMR: -13.6 (*d*, $J = 35.7$); -14.5 (*d*, $J = 35.7$).


b) 383 mg (0.377 mmols) of the product are stirred for 18 hours in 10 ml of ethanol, 1 ml of water and 200 mg of potassium hydroxide. Then, the mixture is concentrated on a rotary evaporator and extracted by shaking in 10 ml of methylene chloride and 10 ml of 2N HCl, and the HCl phase is extracted several times with methylene chloride. The methylene phases are combined, washed with 0.2N HCl and then with water, dried with sodium sulphate and concentrated on a rotary evaporator. 333 mg of product are obtained (white solid, yield 95%).

$^1\text{H-NMR}$: 7.40-6.9 (*m*, 26 arom. H); 4.95 (*s*, 1H, NH), 4.2 (*d*, $J = 14$, 1 H, CH_2O), 3.77 (*d*, $J = 14$, 1 H CH_2O); 3.66 (*t*, 6H, $\text{O-CH}_2\text{CH}_2$), 3.12 (*s*, 6H, $\text{C-CH}_2\text{-O}$), 2.51 (*t*, 6H, $\text{O-CH}_2\text{CH}_2$), 1.38 (*s*, 3 H, Ph-CH_3).

$^{31}\text{P}\{^1\text{H}\}$ -NMR: -13.7 (*d*, $J = 35.2$); -14.5 (*d*, $J = 35.2$).

Example C13: Preparation of a diphosphine of formula

[0148]



a) A solution of 172 mg (0.31 mmols) of (S)-2,2'-dihydroxy-6,6'-diphenylphosphino-diphenyl in 5 ml of methylene chloride is mixed with 0.3 ml of 30% hydrogen peroxide and the mixture is stirred vigorously over night at room temperature. A white precipitate is formed. This poorly soluble precipitate is filtered, washed with water, ethyl acetate and methylene chloride and dried under a vacuum. The yield is practically quantitative. In ^{31}P -NMR (suspension in CD_3OD / DMSO), the poorly soluble white product has one singlet at $\delta + 33.88$ and contains no more educt.

b) 119 mg (0.44 mmols) of N-(3-bromopropyl)phthalimide are added to 104 mg (0.107 mmols) of the accordingly obtained (S)-2,2'-dihydroxy-6,6'-diphenyloxyphosphinodiphenyl and 100 mg of potassium carbonate in 2 ml of N, N-dimethylformamide, and the mixture is stirred over night at 83°C. After cooling, 25 ml of water are added, whereby the product precipitates in pure form practically quantitatively. It is filtered, washed with water and dried under vacuum at 40°C.

^{31}P -NMR (CDCl_3): +29.9 ppm (*s*).

$^1\text{H-NMR}$ (CDCl_3) characteristic signals: 3.7 (*m*, 2H, 2 x $\text{O-CHH}'$) and 3.33 each (*m*, 2H, 2 x $\text{O-CHH}'$), 3.33 (*t*, 4H, 2 x $\text{CH}_2\text{-N}$), 2.5 (*m*, 4H, 2 x $\text{C-CH}_2\text{-C}$).

[0149] The white powder may be reacted to the title compound without further purification, by releasing the amino groups with hydrazine hydrate and reducing the phosphine oxide with trichlorosilane.

D) Preparation of polymers with immobilised diphenyldiphosphines

Example D1: Compound C4 on silica gel

[0150] 2.458 g silica gel (Grace 332; 35-70 mm; Code: 91146025.01) are dried for three hours at 130 °C under a high vacuum and cooled again to room temperature. Then, a solution of 240 mg of compound C4 (2.8 mmols) in 11.5 ml of absolute toluene is added and stirring is carried out over night at 80-90°C. The mixture is cooled to room temperature and the toluene decanted off. The silica gel is then washed five times with 15 ml of degassed methanol and twice with 15 ml of absolute Et_2O , and dried over night under a high vacuum at 40°C. Yield: 2.46 g. analysis: 0.26 %P corresponding to 0.042 mmols diphosphine/g silica gel.

Example D2: Compound C4 on silica gel

[0151] 2.91 g silica gel (Grace 332; 35-70mm; Code: 91146025.01) are dried for three hours at 130°C under a high vacuum and cooled again to room temperature. Then, a solution of 300 mg of compound C4 (0.35 mmols) in 14.5 ml of absolute toluene and 10 ml of methanesulphonic acid is added. The mixture is stirred over night at 90-110°C. After cooling to room temperature, the toluene is decanted off, and the silica gel is washed five times with 15 ml of degassed methanol and twice with absolute Et₂O. The silica gel is then dried over night under a high vacuum at 40°C. Yield: 3.04 g. analysis: P content: 0.39 % corresponding to a density of 0.064 mmols diphosphine/g silica gel.

Example D3: Compound C3 on poly-(bisphenol-A-bisglycidylether) (phenoxy resin)

[0152] 0.4 g of polyphenoxy resin with an average molecular weight of 13,000 (1.41 mmols) are dissolved at 40 °C in 10 ml of absolute CH₂Cl₂. Then, 3.24 ml of 2,4-tolylene-diisocyanate (22.5 mmols) and 20 ml of triethylamine are added. The solution is then stirred under reflux for 3.5 hours. After cooling to room temperature, the polymer is precipitated with 25 ml of absolute pentane and decanted off. The polymer is then washed three times (dissolving in 10 ml of absolute CH₂Cl₂, precipitating with 25 ml of absolute pentane and filtering). Afterwards, the polymer is almost completely dissolved again in 10 ml of CH₂Cl₂. To this are added 220 mg of compound C3 (0.36 mmols) and 10 ml of dibutyltin dilaurate. The solution is stirred under reflux for 18 hours and cooled to room temperature. 6 ml of degassed ethanol and 10 ml of dibutyltin are added and stirring is again effected under reflux for 18 hours. After cooling to room temperature, the polymer is precipitated with 50 ml of absolute pentane and 10 ml of absolute Et₂O and filtered off. The polymer is washed three times (dissolving in 10 ml of CH₂Cl₂, precipitating with 25 ml of absolute pentane and 20 ml of absolute Et₂O and filtering). The polymer is then dried over night under a high vacuum. Yield: 510 mg. Analysis: P: 2.14 % corresponding to a density of 0.34 mmols ligand/g polymer.

Example D4: Compound C3 with 2,4-tolylene-diisocyanate on aminomethylated polystyrene

[0153] 660 mg of aminomethylated polystyrene (NH₂: 0.56 mmols/g polymer) (0.37 mmols) are mixed with 25 ml of absolute CH₂Cl₂. Then, 0.85 ml of 2,4-tolylene-diisocyanate (5.9 mmols) are added dropwise. The mixture is stirred for 3 hours. The solution is filtered and the polymer washed four times with 20 ml of absolute CH₂Cl₂. The polymer is mixed with 20 ml of CH₂Cl₂ and then with 225 mg of compound C3 (0.37 mmols), and then 10 ml of dibutyltin dilaurate are added. The mixture is stirred under reflux for 20 hours and then mixed with 8 ml of degassed EtOH, and with a further 10 ml of dibutyltin dilaurate. The mixture is stirred under reflux for a further 18 hours. After cooling to room temperature, the solvent is filtered off and the polymer washed four times with 25 ml of absolute CH₂Cl₂. The polymer is then dried under a high vacuum. Yield: 754 mg.

Analysis: 0.1% P, corresponding to a density of 0.016 mmols ligand/g polymer.

Example D5: Compound C11 on silica gel

[0154] 2.65 g silica gel (Grace 332; 35-70 mm; Code: 91146025.01) are dried for 4.5 hours at 130°C under a high vacuum and cooled again to room temperature. Then, a solution of 260 mg of compound C11 (0.32 mmols) in 13 ml of absolute toluene and 10 ml of methanesulphonic acid is added. The mixture is stirred over night at 90-105 °C. After cooling to room temperature, the toluene is decanted off, and the silica gel is washed five times with 15 ml of degassed MeOH and twice with absolute Et₂O. The silica gel is then dried over night under a high vacuum at 40°C. Yield: 2.76 g. Analysis: P content: 0.39 % corresponding to a density of 0.0629 mmols ligand/g silica gel.

Example D6: Compound C6(c) with hexylene-1,6-diisocyanate on aminomethylated polystyrene

[0155] 700 mg of aminomethylated polystyrene (NH₂: 0.56 mmols/g polymer) (0.392 mmols) are mixed with 20 ml of absolute CH₂Cl₂. Then, 2.52 ml of hexamethylene-1,6-diisocyanate (16 mmols) are added dropwise. The mixture is stirred for 2.5 hours. The solution is filtered and the polymer washed four times with 20 ml of absolute CH₂Cl₂. The polymer is then mixed with 20 ml of CH₂Cl₂, and 200 mg of compound C6(c) (0.35 mmols) and 10 ml of dibutyltin dilaurate are added. The mixture is stirred under reflux for 22 hours and then mixed with 8 ml of degassed ethanol (EtOH), and 10 ml of dibutyltin dilaurate. The mixture is stirred under reflux for a further 18 hours. After cooling to room temperature, the solvent is filtered off and the polymer washed four times with 25 ml of absolute CH₂Cl₂. The polymer is then dried under a high vacuum. Yield: 808 mg.

Analysis: 0.3% P corresponding to a density of 0.0484 mmols ligand/g polymer.

E) Hydrogenation

Examples E1-E14: Hydrogenation of acetamidocinnamic acid methyl ester

- [0156] Hydrogenation is carried out as follows: 0.0125 mmols of $[\text{Rh}(\text{NBD})_2]\text{BF}_4$ and 0.015 mmols of polymer are placed under argon in a flask having a magnetic stirrer by repeatedly applying a vacuum and rinsing with argon. Then, 2 ml of degassed methanol are added, the mixture is stirred slowly for 10 minutes (if there are insoluble immobilised ligands, the solution loses its colour and the carrier becomes orange) and 8 ml of a degassed methanolic solution of 2.5 mmols of acetamidocinnamic acid methyl ester are added. Then, without stirring, the argon is replaced by hydrogen (normal pressure) by applying a vacuum and rinsing with hydrogen, and hydrogenation is started by switching on the stirrer. The progress of the reaction can be followed by the consumption of hydrogen (pressure drop in the hydrogen reservoir). The results are listed in table 1.

Table 1:

No.	ligand/ polymer ligand	substrate to catalyst	conversion %	reaction time (min.)	optical yield (ee)	remarks
--	Z_1	200	100	60	29 (S)	
--	Z_2	200	100	50	27(R)	
E1	C3	200	100	55	41(S)	
E2	C4	200	100	60	39(S)	
E3	C6(C)	200	100	70	38(R)	
E4	C11	200	100	90	36(R)	
E5	D1	200	100	70	44(S)	
E6	D1	200	100	120	43(S)	reused
E7	D1	1000	93	900	39(S)	
E8	D2	200	100	60	45(S)	
E9	D3	200	98	90	48(R)	catalyst insoluble
E10	D3	200	42	125	46(R)	reused
E11	D3	200	100	60	41(S)	in methanol/THF with methanesulphonic acid
E12	D4	200	100	80	40(R)	
E13	D4	200	100	110	40(R)	reused
E14	D5	400	50	840	39(R)	in methanol/THF with methanesulphonic acid
Z_1 : (S)-6,6'-dimethoxy-diphen-2,2'-yl-bis-(diphenylphosphine)						
Z_2 : (S)-6,6'-dimethyl-diphen-2,2'-yl-bis-(diphenylphosphine)						

Examples E15-E34: Hydrogenation of phenylglyoxalic acid methyl ester to S-mandelic acid methyl ester

- [0157] Hydrogenation is carried out as follows: 2 ml of degassed acetone are added under argon to 0.013 mmols of $(\text{cyclooctadiene})\text{Ru}(\text{2-methylallyl})_2$ and 0.0143 mmols of immobilised ligand D1. Then, the indicated amounts of HBr or methanesulphonic acid or LiBr are added, stirring is effected for 30 minutes, and finally the acetone is drawn off under vacuum. Then, a degassed solution of 5 mmols of phenylglyoxalic acid methyl ester in 5 ml of methanol is added, the reaction mixture is transferred by a cannula under a countercurrent of argon into a 50 ml autoclave, and hydrogenated at 80 bars hydrogen pressure at 40°C.


[0158] The immobilised catalyst is reused several times. Separation takes place each time by centrifuging. The results are listed in table 2. In examples E24-34, 4.6 equivalents of LiBr/Ru are additionally used. The acid addition is given in equivalents per Ru. MS signifies methanesulphonic acid.

Table 2:

No.	acid addition	LiBr addition	substrate to catalyst	conversion (%)	ee (%)	reaction time (min.)	remarks
E15	3.4 HBr	-	384	98.5	86.1	1320	fresh catalyst
E16	-	-	384	93.6	75.7	1100	1st reuse
E17	7.4 HBr	-	384	99	87.1	1080	fresh catalyst
E18	7.4 HBr	-	384	99	89.1	1100	1st reuse
E19	7.4 HBr	-	384	99	88.7	960	2nd reuse
E20	7.4 HBr	-	384	97	88.6	960	3rd reuse
E21	7.4 HBr	-	384	99	90	3780	4th reuse
E22	4.5 HBr	-	1000	47	79	980	fresh catalyst
E23	4.5 HBr	-	1000	65	81.8	1200	1st reuse
E24	2.3 MS	4.6 LiBr	384	79	81.9	870	fresh catalyst
E25	2.3 MS	4.6 LiBr	384	95	86.7	1050	1st reuse
E26	4.6 MS	4.6 LiBr	384	99	90.7	960	2nd reuse
E27	2.3 MS	4.6 LiBr	384	99	86.7	960	3rd reuse
E28	4.6 MS	4.6 LiBr	384	99	87.7	900	4th reuse
E29	4.6 MS	4.6 LiBr	1000	02	85.8	1440	5th reuse
E30	4.6 MS	4.6 LiBr	384	97	86.3	1200	6th reuse
E31	4.6 MS	4.6 LiBr	384	97	87.1	1140	7th reuse
E32	4.6 MS	4.6 LiBr	1000	76	83.7	1200	8th reuse
E33	4.6 MS	4.6 LiBr	1000	57	81.3	900	9th reuse
E34	4.6 MS	4.6 LiBr	1000	35	74.1	1200	10th reuse

Claims

1. Compounds of formula 1,




(I),

wherein

R_1 is methyl chloride, methyl bromide or methyl iodide, R_2 is C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy or has the same significance as R_1 , and R_3 is Br, I or $-NH_2$.

2. Compounds of formula II,




(II),

wherein

R₃ is Br, I or -NH₂,R₄ is hydroxymethyl, aminomethyl, hydroxy-, amino- or cyano-C₂-C₈-alkoxy, hydroxy-, amino- or cyano-C₂-C₈-alkoxymethyl, or hydroxy-, amino- or cyano-C₂-C₈-alkylaminomethyl, and R₅ has the same significance as R₄, or R₅ is C₁-C₄-alkyl or C₁-C₄-alkoxy, orR₄ and R₅ together are HOCH(CH₂-O-)₂, H₂NCH(CH₂-O-)₂, or hydroxy-, amino- or cyano-C₂-C₈-alkyloCH(CH₂-O-)₂, except 2,2'-Dihydroxymethyl-6,6'-dibromodiphenyl and 2,2'-Dihydroxymethyl-6,6'-diaminodiphenyl.

3. Compounds of formula III,



(III),

wherein

R₆ and R₇ signify identical or different secondary phosphino,R₈ is -CH₂-OH, -CH₂-NH₂, -CH₂-O-B-(FU)_p, -CH₂-NR'¹-B-(FU)_p, or -O-B-(FU)_p,R₉ has the same significance as R₈ or is C₁-C₄-alkyl or C₁-C₄-alkoxy, orR₈ and R₉ together signify HOCH(CH₂-O-)₂, H₂NCH(CH₂-O-)₂, (FU)_p-B-OCH(CH₂-O-)₂ or (FU)_p-B-R'¹NCH(CH₂-O-)₂,R'¹ is H or C₁-C₄ alkyl;


B is a bridging group,

FU is a functional group,

p is a number from 1 to 6, and




NH₂ groups are present as such or as masked isocyanate groups.

4. Compounds according to claim 3, in which the secondary phosphino group corresponds to formula -PR₁₀R₁₁, wherein R₁₀ and R₁₁, Independently of one another, are C₁-C₁₂-alkyl, C₅-C₁₂-cycloalkyl, phenyl, C₅-C₁₂-cycloalkyl substituted by C₁-C₄-alkyl or C₁-C₄-alkoxy, or phenyl mono- to trisubstituted by C₁-C₄-alkyl, C₁-C₄-alkoxy, -SiR₁₂R₁₃R₁₄, halogen, -SO₃M, -CO₂M, -PO₃M, -NR₁₅R₁₆, -[⁺NR₁₅R₁₆R₁₇]X⁻ or C₁-C₅-fluoroalkyl; R₁₀ and R₁₁ together are tetra- or pentamethylene either unsubstituted or mono- to trisubstituted by C₁-C₄-alkyl, C₁-C₄-alkoxy, -SiR₁₂R₁₃R₁₄, halogen, -SO₃M, -CO₂M, -PO₃M, -NR₁₅R₁₆, -[⁺NR₁₅R₁₆R₁₇]X⁻ or C₁-C₅-fluoroalkyl, or the group -PR₁₀R₁₁ represents a radical of formulae



10 and R_{12} , R_{13} and R_{14} independently of one another, are C_1 - C_{12} -alkyl or phenyl
 R_{15} and R_{16} , independently of one another, are H, C_1 - C_{12} -alkyl or phenyl, or R_{15} and R_{16} together are tetramethylene, pentamethylene or 3-oxa-1,5-pentylene;
 R_{17} is H or C_1 - C_4 -alkyl;
M is H or an alkali metal;
15 X is the anion of a monobasic acid;
halogen is fluorine, chlorine, bromine or iodine.

5. Metal complexes of formulae V, Va and Vb of d-8 metals with the compounds of formula III,



whereby R_6 , R_7 , R_8 and R_9 have the above-mentioned significances and preferences;

Y denotes two monoolefin ligands or one diene ligand;


Me signifies a d-8 metal selected from the group Ir and Rh;

D is -Cl, -Br or -I; and

E is the anion of an oxyacid or complex acid;


X_2 and X'_2 are identical or different and have the significance of D or E, or X_2 and X'_2 are allyl or 2-methylallyl, or X_2 has the significance of D and E and X'_2 is hydride.

6. Solid inorganic carrier, which has diphosphine ligands of formula IIIa that are bonded at the surface by one or two silyl groups of the radical of formula




(IIIa),

wherein the group B-FU in radicals R₈ and R₉ is a radical of formula



wherein X₃ signifies -O-, -NH- or NH(C₁-C₄-alkyl), R₂₇ is C₁-C₁₂-alkylene, R₂₆ is C₁-C₁₂ alkyl, R₂₅ is C₁-C₄-alkyl or OR₂₆ and n is 0, 1 or 2.



7. Inorganic or organic polymeric carriers, to which diphosphines of formula III




(III)

are bonded, which are **characterised in that** they are bonded to the inorganic or polymeric organic carrier by the functional group FU, whereby the radicals FU, R₆, R₇, R₈ and R₉ have the significances given in claim 3.

8. d-8 metal complexes of inorganic or organic polymeric carriers, to which diphenyldiphosphines of formula VII, VIIa or VIIb





are bonded by at least one HO-, H₂N- group or functional group FU, whereby the radicals R₆, R₇, R₈, R₉, Me, E, Y, X₂ and X₂' and the carrier have the significances given in claim 5.

9. Diphenyldiphosphine ligands of formula III according to claim 3 and their d-8 metal complexes of formulae V, Va and Vb according to claim 5, which contain solubility-enhancing or adsorption-facilitating groups bonded to at least one HO-, H₂N- or functional group FU, and which have a molecular weight of less than 5000 daltons, and which can be separated by extraction with immiscible liquids or by adsorption on a carrier.

10. Process for the asymmetric hydrogenation of compounds with carbon double bonds or carbon/hetero atom double bonds, whereby the compounds are reacted with hydrogen at a temperature of -20 to 80°C and at a hydrogen pressure of 10⁵ to 2x10⁷ Pa in the presence of catalytic amounts of d-8 metal complexes according to claims 5, 8 or 9.

Patentansprüche

1. Verbindungen der Formel I,



worin

R₁ Chlormethyl, Brommethyl oder Iodmethyl ist, R₂ C₁-C₄-Alkyl oder C₁-C₄-Alkoxy bedeutet oder die gleiche Bedeutung wie R₁ hat, und R₃ für Br, I oder -NH₂ steht.

2. Verbindungen der Formel II,



worin


R_3 für Br, I oder $-NH_2$ steht,

R_4 Hydroxymethyl, Aminomethyl, Hydroxy-, Amino- oder Cyano- C_2-C_8 -alkoxy, Hydroxy-, Amino- oder Cyano- C_2-C_8 -alkoxymethyl, oder Hydroxy-, Amino- oder Cyano- C_2-C_8 -alkylaminomethyl bedeutet, und R_5 die gleiche Bedeutung wie R_4 hat, oder R_5 C_1-C_4 -Alkyl oder C_1-C_4 -Alkoxy darstellt, oder

R_4 und R_5 zusammen $HOCH(CH_2-O-)_2$, $H_2NCH(CH_2-O-)_2$, oder

Hydroxy-, Amino- oder Cyano- C_2-C_8 -alkyl $OCH(CH_2-O-)_2$ sind, ausgenommen 2,2'-Bis-(hydroxymethyl)-6,6'-dibrombiphenyl und 2,2'-Bis-(hydroxymethyl)-6,6'-diaminobiphenyl.

3. Verbindungen der Formel III,



(III),

worin

R_6 und R_7 gleiches oder verschiedenes Sekundärphosphino bedeuten,

R_8 $-CH_2-OH$, $-CH_2-NH_2$, $-CH_2-O-B-(FU)_p$, $-CH_2-NR'-B-(FU)_p$, oder $-O-B-(FU)_p$ darstellt,

R_9 die gleiche Bedeutung wie R_8 hat oder für C_1-C_4 -Alkyl oder C_1-C_4 -Alkoxy steht, oder

R_8 und R_9 zusammen $HOCH(CH_2-O-)_2$, $H_2NCH(CH_2-O-)_2$, $(FU)_p-B-OCH(CH_2-O-)_2$ oder $(FU)_p-B-R'NCH(CH_2-O-)_2$ bedeuten,

R' für H oder C_1-C_4 -Alkyl steht,


B eine Brückengruppe ist,

FU eine funktionelle Gruppe bedeutet,

p für eine Zahl von 1 bis 6 steht, und

NH_2 -Gruppen als solche oder als verkappte Isocyanatgruppen vorliegen.

4. Verbindungen gemäss Anspruch 3, **dadurch gekennzeichnet, dass** die Sekundärphosphinogruppe der Formel $-PR_{10}R_{11}$ entspricht, worin R_{10} und R_{11} unabhängig voneinander C_1-C_{12} -Alkyl, C_5-C_{12} -Cycloalkyl, Phenyl, mit C_1-C_4 -Alkyl oder C_1-C_4 -Alkoxy substituiertes C_5-C_{12} -Cycloalkyl oder mit ein bis drei C_1-C_4 -Alkyl, C_1-C_4 -Alkoxy, $-SiR_{12}R_{13}R_{14}$, Halogen, $-SO_3M$, $-CO_2M$, $-PO_3M$, $-NR_{15}R_{16}$, $-[{}^+NR_{15}R_{16}R_{17}]X^-$ oder C_1-C_5 -Fluoralkyl substituiertes Phenyl; R_{10} und R_{11} zusammen unsubstituiertes oder mit ein bis drei C_1-C_4 -Alkyl, C_1-C_4 -Alkoxy, $-SiR_{12}R_{13}R_{14}$, Halogen, $-SO_3M$, $-CO_2M$, $-PO_3M$, $-NR_{15}R_{16}$, $-[{}^+NR_{15}R_{16}R_{17}]X^-$ oder C_1-C_5 -Fluoralkyl substituiertes Tetra- oder Pentamethylen darstellt, oder die Gruppe $-PR_{10}R_{11}$ einen Rest der Formeln darstellt,



10 und R_{12} , R_{13} und R_{14} unabhängig voneinander C_1 - C_{12} -Alkyl oder Phenyl

R_{15} und R_{16} unabhängig voneinander H, C_1 - C_{12} -Alkyl oder Phenyl oder R_{15} und R_{16} zusammen Tetramethylen, Pentamethylen oder 3-Oxa-1,5-Pentyl;




15 R_{17} H oder C_1 - C_4 -Alkyl;

M H oder ein Alkalimetall;

X^- das Anion einer einbasischen Säure;

20 Halogen Fluor, Chlor, Brom oder Iod bedeuten.

5. Metallkomplexe der Formel V, Va und Vb von d-8-Metallen mit den Verbindungen der Formeln III,



45 wobei R_6 , R_7 , R_8 und R_9 die vorstehend angegebenen Bedeutungen und Bevorzungen haben;

Y für zwei Monoolefinliganden oder einen Dienliganden steht;


50 Me ein d-8-Metall ausgewählt aus der Gruppe Ir und Rh bedeutet;

D -Cl, -Br oder -I darstellt; und

E das Anion einer Sauerstoffsäure oder Komplexsäure ist,


55 X_2 und X_2' gleich oder verschieden sind und die Bedeutung von D oder E haben, oder X_2 und X_2' für Allyl oder 2-Methylallyl stehen, oder X_2 die Bedeutung von D und E hat und X_2' für Hydrid steht.

6. Festes anorganisches Trägermaterial, **dadurch gekennzeichnet, dass** es über ein oder zwei Silylgruppen an der Oberfläche gebundene Diphosphinliganden der Formel aufweist,




(IIIa),

worin die Gruppe B-FU in den Resten R_8 und R_9 einen Rest der Formel



darstellt, worin X_3 -O-, -NH- oder $NH(C_1-C_4\text{-Alkyl})$ bedeutet, R_{27} C_1-C_{12} -Alkyl, R_{26} C_1-C_{12} Alkyl, R_{25} C_1-C_4 -Alkyl oder OR_{26} bedeuten und n 0, 1 oder 2 ist.



7. Anorganische oder organische polymere Trägermaterialien, an die Diphosphine der Formel III




(III)

gebunden sind, die **dadurch gekennzeichnet sind, dass** sie über die funktionelle Gruppe FU an das anorganische oder polymere organische Trägermaterial gebunden sind, wobei die Reste FU, R_6 , R_7 , R_8 und R_9 die in Anspruch 3 angegebenen Bedeutungen haben.

8. d-8-Metallkomplexe von anorganischen oder organischen polymeren Trägermaterialien, an die Diphenyldiphosphine der Formel VII, VIIa oder VIIb






(VIIb),

über wenigstens eine HO-, H₂N-Gruppe oder funktionelle Gruppe FU an das anorganische oder organische polymere Trägermaterial gebunden sind, wobei die Reste R₆, R₇, R₈, R₉, Me, E, Y, X₂ und X'₂ und das Trägermaterial die in Anspruch 5 angegebenen Bedeutungen haben.

9. Diphenyldiphosphinliganden der Formel III gemäß Anspruch 3 und deren d-8 Metallkomplexe der Formeln V, Va und Vb gemäß Anspruch 5, welche an wenigstens eine HO-, H₂N- oder funktionelle Gruppe FU löslichkeitsvermittelnde oder adsorptionsvermittelnde Gruppen gebunden enthalten, und die ein Molekulargewicht von weniger als 5000 Dalton, welche durch Extraktion mit nicht miteinander mischbaren Flüssigkeiten oder durch Adsorption an einem Trägermaterial aus abtrennbar sind.
10. Verfahren zur asymmetrischen Hydrierung von Verbindungen mit Kohlenstoff- oder Kohlenstoff-Heteroatomdoppelbindungen, **dadurch gekennzeichnet, dass** man die Verbindungen bei einer Temperatur von -20 bis 80 °C und einem Wasserstoffdruck von 10⁵ bis 2x10⁷ Pa in Gegenwart katalytischer Mengen von d-8 Metallkomplexen gemäß den Ansprüchen 5, 8 oder 9 mit Wasserstoff umsetzt.

Revendications

1. Composés de formule 1,




(I),


dans laquelle

R₁ est le chlorure de méthyle, le bromure de méthyle ou l'iodure de méthyle, R₂ est un groupe alkyle en C₁ à C₄ ou un groupe alcoxy en C₁ à C₄ ou possède les mêmes significations que R₁, et R₃ est Br, I ou -NH₂.

2. Composés de formule II,



(II),




(II),

dans laquelle

R_3 est Br, I ou $-NH_2$,

R_4 est un groupe hydroxyméthyle, aminométhyle, hydroxy-, amino- ou cyano-(alcoxy en C_2 à C_8), hydroxy-, amino- ou cyano-(alcoxy en C_2 à C_8)-méthyle, ou hydroxy-, amino- ou cano-(alkyle en C_2 à C_8)-aminométhyle, et R_5 possède la même signification que R_4 , ou R_5 est un groupe alkyle en C_1 à C_4 ou alcoxy en C_1 à C_4 , ou R_4 et R_5 , pris conjointement, sont $HOCH(CH_2-O)_2$, $H_2NCH(CH_2-O)_2$, ou un groupe hydroxy-, amino- ou cyano-(alkyle en C_2 à C_8)- $OCH(CH_2-O)_2$, sauf le 2,2'-dihydroxyméthyl-6,6'-dibromodiphényle et le 2,2'-dihydroxyméthyl-6,5'-diaminodiphényle.

3. Composés de formule III,



(III),

dans laquelle

R_6 et R_7 représentent un groupe phosphino secondaire identique ou différent,

R_8 représente $-CH_2-OH$, $-CH_2-NH_2$, $-CH_2-O-B-(FU)_p$, $-CH_2-NR'-B-(FU)_p$ ou $-O-B-(FU)_p$,

R_9 possède la même signification que R_8 ou est un groupe alkyle en C_1 à C_4 ou alcoxy en C_1 à C_4 , ou

R_8 et R_9 , pris conjointement, représentent $HOCH(CH_2-O)_2$, $H_2NCH(CH_2-O)_2$, $(FU)_p-B-OCH(CH_2-O)_2$ ou $(FU)_p-B-R'NCH(CH_2-O)_2$,


R' est un atome H ou un groupe alkyle en C_1 à C_4 ,

B est un groupe pontant,

FU est un groupe fonctionnel,


p est un nombre compris entre 1 et 6, et les radicaux NH_2 sont présents tels quels ou en tant que groupes isocyanates masqués.

4. Composés selon la revendication 3, dans lesquels le groupe phosphino secondaire correspond à la formule $-PR_{10}R_{11}$, dans laquelle R_{10} et R_{11} , indépendamment l'un de l'autre, sont un groupe alkyle en C_1 à C_{12} , cycloalkyle en C_5 à C_{12} , phényle, cycloalkyle en C_5 à C_{12} substitué par un groupe alkyle en C_1 à C_4 ou alcoxy en C_1 à C_4 , ou phényle mono- à trisubstitué par un groupe alkyle en C_1 à C_4 , alcoxy en C_1 à C_4 , $-SiR_{12}R_{13}R_{14}$, halogéno, $-SO_3M$, $-CO_2M$, $-PO_3M$, $-NR_{15}R_{16}$, $-[+NR_{15}R_{16}R_{17}]X^-$ ou fluoroalkyle en C_1 à C_5 ; R_{10} et R_{11} , pris conjointement, sont le tétra- ou pentaméthylène, soit non substitué, soit mono- à trisubstitué par un groupe alkyle en C_1 à C_4 , alcoxy en C_1 à C_4 , $-SiR_{12}R_{13}R_{14}$, halogéno, $-SO_3M$, $-CO_2M$, $-PO_3M$, $-NR_{15}R_{16}$, $-[+NR_{15}R_{16}R_{17}]X^-$ ou fluoroalkyle en C_1 à C_5 ou le groupe $-PR_{10}R_{11}$ représente un radical de formules :



et R_{12} , R_{13} et R_{14} , indépendamment les uns des autres, sont un groupe alkyle en C_1 à C_{12} ou phényle, R_{15} et R_{16} , indépendamment l'un de l'autre, représentent un atome H, un groupe alkyle en C_1 à C_{12} ou phényle, ou R_{15} et R_{16} , pris conjointement, sont le tétraméthylène, le pentaméthylène ou le 3-oxa-1,5-pentylène ; R_{17} est un atome H ou un groupe alkyle en C_1 à C_4 ; M est un atome H ou de métal alcalin ; X est l'anion d'un acide monobasique ; halogéno représente un groupe fluoro, chloro, bromo ou iodo.

5. Complexes de métaux de formules V, Va et Vb des métaux d-8 avec les composés de formule III,




dans laquelle R_6 , R_7 , R_8 et R_9 possèdent les significations et les préférences mentionnées ci-dessus ;

Y représente deux ligands monooléfiniques ou un ligand diène ;
 M signifie un métal d-8 choisi dans le groupe formé par Ir et Rh ;
 D est -Cl, -Br ou -I ; et
 E est l'anion d'un oxyacide ou d'un acide complexe ;


X_2 et X_2' sont identiques ou différents et possèdent la signification de D ou E, ou X_2 et X_2' sont un groupe allyle ou 2-méthylallyle, ou X_2 , possède la signification de D et E et X_2' est un hydrure.

6. Support inorganique solide, qui possède des ligands diphosphine de formule IIIa qui sont reliés à la surface par un ou deux groupes silyle du radical de formule :




(IIIa),

dans laquelle le groupe B-FU dans les radicaux R_8 et R_9 est un radical de formule :



dans laquelle X_3 signifie $-O-$, $-NH-$ ou NH (alkyle en C_1 à C_4), R_{27} est un groupe alkylène en C_1 à C_{12} , R_{26} est un groupe alkyle en C_1 à C_{12} , R_{25} est un groupe alkyle en C_1 à C_4 ou OR_{26} et n vaut 0, 1 ou 2.



7. Supports polymères inorganiques ou organiques, auxquels les diphosphines de formule III




(III)

sont reliées, qui sont caractérisés en ce qu'ils sont reliés au support inorganique ou polymère organique par le groupe fonctionnel FU, dans laquelle les radicaux FU, R_6 , R_7 , R_8 et R_9 possèdent les significations données dans la revendication 3.

8. Complexes de métaux d-8 de supports polymères inorganiques ou organiques, auxquels les diphényldiphosphines de formule VII, VIIa ou VIIb





sont reliées par au moins un groupe HO-, H₂N- ou un groupe fonctionnel FU, dans lesquelles les radicaux R₆, R₇, R₈, R₉, Me, E, Y, X₂ et X₂' et, le support possèdent les significations données dans la revendication 5.

- 15 9. Ligands diphényldiphosphines de formule III, selon la revendication 3 et leurs complexes de métaux d-8 de formules V, Va et Vb selon la revendication 5, qui contiennent des groupes favorisant la solubilité ou facilitant l'adsorption reliés à au moins un groupe HO-, H₂N- ou un groupe fonctionnel FU, et qui ont une masse moléculaire inférieure à 5 000 daltons, et qui sont séparés par extraction avec des liquides immiscibles ou par adsorption sur un support.
- 20 10. Procédé pour l'hydrogénation asymétrique de composés avec des doubles liaisons carbone ou des doubles liaisons carbone/hétéroatome, dans lequel les composés sont mis à réagir avec de l'hydrogène à une température comprise entre -20°C et 80°C et à une pression d'hydrogène de 10⁵ à 2x10⁷ Pa en présence de quantités catalytiques de complexes de métaux d-8 selon les revendications 5, 8 ou 9.
- 25
- 30
- 35
- 40
- 45
- 50
- 55